

## Surveillance and outbreak reports

### DEMOGRAPHIC FEATURES AND TRENDS IN TUBERCULOSIS CASES IN THE EUROPEAN REGION, 1995-2005

D Falzon (d.falzon@invs.sante.fr)<sup>1</sup>, D van Caunteren<sup>1,2</sup>

1. Département des maladies infectieuses, Institut de veille sanitaire (Department of infectious diseases, French Institute for Public Health Surveillance), Saint-Maurice, France

2. Programme de formation à l'épidémiologie de terrain (PROFET), Département des maladies infectieuses, Institut de veille sanitaire (Department of infectious diseases, French Institute for Public Health)

In 2005, 426,457 tuberculosis (TB) cases were notified in the World Health Organization (WHO) European Region, with a wide variation and an incremental west-to-east gradient in notification rates also reflected in TB mortality rates. In the enlarged European Union ('EU-27') and other western countries - where 19% of cases were of foreign origin in 2005 (>50% in 13 countries) - overall TB notification rates decreased by 2.4% yearly between 2000 and 2005, compared to 1.6% in 1995-2000, reflecting a declining incidence in all age groups and in most countries. Half the cases notified by the 12 ex-republics of the former Soviet Union in the East in 2005 were reported by the Russian Federation. In the East, the mean annual increase in 1995-2000 (10.0%) was higher than in 2000-2005 (3.9%), and in recent years the number of new cases stabilised while previously treated cases have increased. Efforts are still needed to improve and standardise TB surveillance across the Region. The collection of additional data on risk factors of TB may be useful for surveillance and control.

#### Introduction

The last years of the 20th century saw the resurgence of TB incidence in different parts of the world. In western countries of the World Health Organization (WHO) European Region where rates had been falling steadily for many years, the rate of decline decreased and, in some countries, a perceptible increase was observed [1]. At that time, countries in the central and eastern part of the Region were experiencing profound economic upheavals which impacted negatively upon their public health [2]. TB notification rates started increasing in the Baltic States (Estonia, Latvia and Lithuania) and in certain central Asian states after the end of the 1980s [3]. This heralded an upturn in TB rates in other former Soviet Union republics, which was sharper and more protracted than that seen further west. The true extent of the epidemic was difficult to assess since case definitions and completeness of reporting differed. In the wake of these developments, a TB surveillance network – EuroTB (<http://www.eurotb.org>) – was established in 1996 through funding by the European Commission to support TB surveillance across the WHO European Region. This article uses data from this network and from the WHO to describe the main demographic features and trends in TB between 1995 and 2005 in the Region.

#### Methods

Until 2007, the coordinating hub of EuroTB, located at the Institut de veille sanitaire in France, collected TB data from European national surveillance authorities in coordination with the WHO. In this article the 53 countries of the WHO European

Region have been grouped into three geographic areas, based on epidemiological and geographic features (Table 1)

- the European Union and West (EU and West), composed of the 27 current Member States of the EU as well as other industrialized countries of Western Europe (Andorra, Iceland, Israel, Monaco, Norway, San Marino, Switzerland);
- the Balkans (Albania, Bosnia and Herzegovina, Croatia, Former Yugoslav Republic of Macedonia, Montenegro, Serbia, Turkey); and
- the East, made up of 12 republics of the former Soviet Union (Armenia, Azerbaijan, Belarus, Georgia, Kazakhstan, Kyrgyzstan, Moldova, Russian Federation, Tajikistan, Turkmenistan, Ukraine, Uzbekistan).

Data used were updated until 31 December 2007. Population estimates used for calculating TB notification rates were obtained from the United Nations Population Division,<sup>4</sup> with the exception of Serbia, which provided its own estimates for 1998-2005. Updates in notification and population data may account for slight differences in values compared to those published earlier. Data about TB deaths were obtained from the WHO Mortality Database website [5].

The definitions and methodology recommended for use in TB surveillance have been described elsewhere [6,7]. TB cases enumerated in this article include both those which were laboratory confirmed as well as others diagnosed only on clinical/radiological grounds. In 2005, 58% of TB cases in the EU and West (country range: 28-100%) were culture positive, as opposed to 34% in the Balkans (24-57%) and 20% in six countries in the East (4-36%). The geographic origin of TB cases was assigned as 'national' or 'foreign' in relation to the country of report on the basis of patients' place of birth or citizenship. Previously treated cases were those who received curative, combination anti-TB chemotherapy for one month or more prior to the current episode.

Rates of notification and mortality are expressed per 100,000 total population and stratified by age group and sex where indicated. No adjustment for reporting completeness was made. In 2006, countries reported completeness of TB notification to be 70-100% in the EU and West (23/34 countries), 90-98% in the Balkans (2/7 countries) and 48-100% in the East (8/12 countries). Methods used to derive estimates of completeness differed between countries and were sometimes not described. The time trend in notification rates was expressed as the mean of the percentage difference in rates (un-rounded) from one year to the next, and was not shown

for Montenegro (data first reported separately for 2005) and for individual countries reporting less than 60 cases in 2005.

The contribution of HIV to TB morbidity was expressed as the proportion of all notified TB cases known to be positive for HIV. The availability of HIV-testing results among TB patients depends on testing policies and the methodology used to collect the test results, which differed between countries. Only deaths coded ICD-9 010-018 (BTL 020-025, 029) or ICD-10 A15-19 were considered in the calculation of TB mortality. Indicators on other clinical characteristics of TB cases, on laboratory confirmation, on anti-TB drug resistance, and on treatment outcome were beyond the focus of this report.

### Results

Of all the 426,457 TB cases reported in the WHO European Region in 2005, 72% were from the East, 12% from 12 countries joining the EU since 2004, 10% from countries in the original 'EU-15' and West, and 6% from the Balkans including Turkey (Table 1). The overall TB notification rate in 2005 was 47/100,000 population, with an incremental gradient when moving from west to east (country range: 4-205/100,000), which was also reflected in TB mortality (0-25/100,000; Table 2).


Overall the number of TB cases reported yearly in the Region has increased markedly since 1995. This increase however was not uniformly distributed over the years or between the countries. Total notification rates have continued to diverge between the EU and West and the East in recent years (Figure 1).

### EU and West

In 2005, the overall notification rate in the EU and West was 18/100,000, with a rate of 10/100,000 or lower in 15 countries and higher than 30/100,000 in Romania (135), the Baltic States (39-75), Bulgaria (43), and Portugal (34). Scandinavian countries


and countries on the Mediterranean littoral had some of the lowest notification rates in the area. The overall rate in the 12 countries joining the EU since 2004 was over four times that in the original 15 Member States. Despite the inclusion of countries with a higher TB incidence, the overall notification rate in the EU and West in 2005 was 10% lower than it was in 2001, reflecting a downward trend in 20 countries. Overall rates in the 'EU15' and West decreased by a similar gradient in 1995-2000 and in 2000-2005. Greece, Ireland and Sweden had a net increase in 2000-2005 after a decline in 1995-2000. The United Kingdom had an increase throughout 1995-2005, particularly between 2003 and 2005. In the Baltic States, rates decreased in 2000-2005

**FIGURE 1**  
TB notification by area and treatment history, WHO European Region\*, 2001-2005



\* excluding countries with missing data for any year: Cyprus, Monaco, San Marino (EU and West); Bosnia and Herzegovina (Balkans); Belarus, Tajikistan, Ukraine (East)

**FIGURE 2**  
Age-group and sex specific TB notification rates, WHO European Region\*, 2005



\* N=170,795; excluding cases without information on age or sex (407) and cases from countries without age-group distribution of TB cases or population: Andorra, Monaco, San Marino (EU and West); Belarus, Kyrgyzstan, Moldova, Russian Federation, Tajikistan, Ukraine (East). Romania excluded as the age-specific rates are very different from the rest of the EU and West.

TABLE 1

## Total tuberculosis notifications and notification rates, WHO European Region, 1995-2005

	Total TB notifications										Total TB notification rates (per 100,000 population)											
	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
<b>'EU-15' (pre-2004)</b>																						
Austria	1,383	1,489	1,398	1,306	1,245	1,225	1,075	1,076	980	1,061	999											
Belgium	1,380	1,352	1,263	1,203	1,270	1,313	1,321	1,294	1,117	1,198	1,107											
Denmark	448	484	554	529	536	548	511	419	393	385	422											
Finland	662	644	573	629	566	537	494	473	412	331	361											
France	8,723	7,656	6,832	6,651	6,674	6,714	6,465	6,322	6,098	5,514	5,374											
Germany <sup>a</sup>	12,198	11,814	11,163	10,440	9,974	9,064	7,539	7,701	7,166	6,542	6,020											
Greece <sup>a</sup>	939	945	767	1,152	952	703	617	582	620	774	769											
Ireland	458	434	416	424	469	403	406	408	407	432	450											
Italy	5,225	5,152	5,176	4,795	4,429	4,759	4,505	4,212	4,518	4,220	4,137											
Luxembourg	32	36	38	44	42	44	32	32	54	31	37											
Netherlands	1,619	1,678	1,486	1,341	1,535	1,404	1,436	1,401	1,321	1,344	1,155											
Portugal	5,577	5,248	5,112	5,260	5,160	4,494	4,399	4,501	4,148	3,854	3,573											
Spain <sup>b</sup>	8,764	8,331	9,347	9,111	8,393	8,395	7,453	7,626	7,467	7,766	7,820											
Sweden	564	493	456	446	493	458	428	407	408	461	559											
United Kingdom	6,161	6,240	6,355	6,176	6,287	6,792	7,017	7,263	7,220	7,609	8,317											
<b>Subtotal 'EU-15'</b>	<b>54,133</b>	<b>51,996</b>	<b>50,936</b>	<b>49,507</b>	<b>48,025</b>	<b>46,853</b>	<b>43,698</b>	<b>43,717</b>	<b>42,329</b>	<b>41,522</b>	<b>41,100</b>											
<b>New EU countries (since 2004)</b>																						
Bulgaria	3,245	3,109	3,437	4,117	3,530	3,349	3,862	3,335	3,263	3,232	3,302											
Cyprus	36	24	47	45	39	33	40	20	35	30	37											
Czech Republic	1,851	1,936	1,834	1,805	1,631	1,442	1,350	1,200	1,162	1,057	1,007											
Estonia <sup>c</sup>	608	683	744	818	754	791	812	713	623	594	519											
Hungary	4,339	4,278	4,240	3,999	3,914	3,598	3,150	2,838	2,582	2,340	1,964											
Latvia <sup>c</sup>	1,541	1,761	2,003	2,182	1,968	2,063	2,082	1,855	1,726	1,610	1,443											
Lithuania <sup>c</sup>	2,362	2,608	2,926	3,016	2,903	2,981	2,989	2,844	2,821	2,514	2,574											
Malta	10	29	11	16	22	18	16	24	7	19	25											
Poland	15,959	15,358	13,967	13,302	12,179	11,477	10,672	10,475	10,124	9,493	9,280											
Romania	23,271	24,113	23,903	25,758	26,870	27,667	30,440	33,595	31,039	31,034	29,289											
Slovakia	1,537	1,499	1,298	1,282	1,218	1,111	1,076	1,053	983	705	760											
Slovenia	525	563	481	449	438	380	371	350	293	263	278											
<b>Subtotal New EU countries</b>	<b>55,284</b>	<b>55,961</b>	<b>54,891</b>	<b>56,789</b>	<b>55,466</b>	<b>54,910</b>	<b>56,860</b>	<b>58,302</b>	<b>54,658</b>	<b>52,891</b>	<b>50,478</b>											
<b>Subtotal all EU ('EU-27')</b>	<b>109,417</b>	<b>107,957</b>	<b>105,827</b>	<b>106,296</b>	<b>103,491</b>	<b>101,763</b>	<b>100,558</b>	<b>102,019</b>	<b>96,987</b>	<b>94,413</b>	<b>91,578</b>											
	23	22	22	22	21	21	21	21	21	21	21											
	39	38	42	51	44	42	42	49	42	42	42											
	5	3	6	6	5	4	4	5	2	4	4											
	18	19	18	18	16	14	14	13	12	11	10											
	42	48	53	59	55	58	60	60	53	46	44											
	42	41	41	39	38	35	31	28	25	23	19											
	62	72	82	90	82	87	88	79	74	70	63											
	65	72	82	85	82	85	86	82	82	82	75											
	3	8	3	4	6	5	4	4	6	2	5											
	41	40	36	35	32	30	28	27	26	25	24											
	103	107	106	115	121	125	138	153	142	143	135											
	29	28	24	24	23	21	20	20	18	13	14											
	27	29	24	23	22	19	19	18	15	13	14											
	52	53	52	54	53	52	54	54	56	53	49											
	23	22	22	22	21	21	21	21	21	20	19											

West, non-EU		17	19	8	9	11	5	5	5	11	7	10
		12	10	17	12	13	13	8	8	5	12	11
Iceland		398	415	422	656	520	591	564	511	529	519	406
Israel <sup>a</sup>		1	0	0	3	0	0	0	0	1	-	-
Monaco		236	217	205	244	273	288	251	337	302	288	-
Norway		2	0	1	0	1	0	1	1	0	-	-
San Marino		830	764	747	749	772	629	611	658	623	593	567
Switzerland		110,896	109,381	107,231	107,970	105,080	103,245	102,039	103,453	98,494	95,846	92,860
Total EU & West		664	707	655	694	765	631	572	612	561	581	540
Balkans		2,132	2,220	2,869	3,071	3,075	2,606	2,551	2,551	1,780 <sup>d</sup>	2,382	2,160
Albania		2,114	2,174	2,054	2,118	1,770	1,630	1,505	1,470	1,493	1,297	1,141
Bosnia and Herzegovina <sup>d</sup>		786	724	693	620	576	668	697	730	697	680	658
Croatia		-	-	-	-	-	-	-	-	-	-	-
Macedonia, Former Yugoslav Rep.		4,169	4,541	4,062	3,028	2,646	2,922	2,888	3,033	2,949	2,824	2,378
Montenegro		23,035	23,533	25,685	25,501	22,088	18,038	18,890	19,028	18,590	19,799	20,535
Serbia <sup>a</sup>		32,900	33,899	36,018	35,032	30,920	26,495	27,103	27,424	26,070	27,563	27,582
Turkey		836	935	1,026	1,455	1,499	1,344	1,401	1,455	1,570	1,701	2,322
Total Balkans		3,306	5,006	4,635	4,350	4,629	5,187	4,923	5,348	3,931	6,501	7,920
East		5,092	5,619	5,985	5,595	7,339	6,084	5,505	5,139	5,963	6,490	6,357
Armenia		-	10,641	8,446	6,302	6,546	6,436	5,876	6,345	5,993	5,967	6,448
Azerbaijan		11,095	13,559	16,109	20,623	25,060	28,265	31,254	32,936	32,169	32,131	31,187
Belarus		3,380	4,086	5,189	5,935	6,501	6,383	6,901	6,794	7,025	6,641	6,765
Georgia		2,753	2,922	2,908	2,891	2,947	2,935	3,820	4,149	5,027	6,008	6,278
Kazakhstan		96,828	110,897	119,123	121,917	135,054	143,801	138,432	134,812	152,244	152,438	156,047
Kyrgyzstan		2,029	1,647	2,143	2,503	2,553	2,779	3,508	4,052	4,883	5,122	7,142
Moldova		2,009	2,149	3,438	3,712	4,092	3,967	4,922	4,635	4,759	4,172	3,291
Russian Federation		21,459	26,834	28,344	31,318	32,879	32,963	36,784	40,175	40,659	38,403	43,367
Tajikistan		9,866	11,919	13,352	13,958	16,959	15,912	18,106	27,009	26,172	25,714	28,891
Turkmenistan		158,653	196,214	210,698	220,559	246,058	256,056	261,432	272,849	290,395	291,288	306,015
Ukraine		302,449	339,494	353,947	363,561	382,058	385,796	390,574	403,726	414,959	414,697	426,457
Uzbekistan		35	39	41	42	44	44	44	44	45	46	47
Total East		26	29	33	33	47	48	44	44	46	48	56
WHO European Region		42	64	58	54	57	64	60	65	66	66	77
		50	55	59	55	73	61	55	52	52	60	65
		-	215	173	131	137	136	126	138	126	131	144
		70	86	104	135	166	189	210	221	214	213	205
		74	88	110	124	133	129	138	134	138	129	130
		63	67	68	68	70	71	93	103	103	126	162
		65	74	80	82	91	98	94	92	105	105	108
		35	28	36	42	42	45	56	64	76	79	109
		48	50	79	85	92	88	108	100	101	88	68
		42	53	56	63	67	67	76	84	85	81	92
		43	51	56	58	70	64	72	106	101	98	109
		57	69	74	78	87	91	93	97	104	104	110
		35	39	41	42	44	44	45	46	47	47	48

Information updated to 31 December 2007. Not including cases from Abkhazia, Greenland (99 cases in 2005), Northern Cyprus and S.Ossetia; neither cases from Kosovo (1 102 cases in 2005) after 1997. Blue figures indicate new cases only.

a. Important changes to reporting system in Germany (2001) and in Greece and Israel (1998)

b. 1995-1996: new respiratory cases only; 1997-2003: all respiratory and meningial cases; 2004-2005: all cases

c. inclusion of retreated TB cases other than relapses from 2001

d. no data from Republika Srpska in 2003 (rate not shown)

e. included cases from Kosovo until 1997 and from Montenegro until 2004

TABLE 2

## Tuberculosis notification and mortality, WHO European Region: selected indicators

Country	Mean annual % change in total notification rate		TB notification, 2005							HIV status of TB cases			TB mortality, 2005 <sup>a</sup>	
	1995-2000	2000-2005	% cases in males	% cases in <15y	% cases in >64y	% cases of foreign origin <sup>b</sup>	% cases treated for TB in past	Cases with HIV infection	% HIV status unknown	N	Rate/100,000 population			
	N	% <sup>c</sup>												
<b>'EU-15' (pre-2004)</b>														
Austria	-2.4	-4.2	66%	4.2%	20.9%	44%	3%	-	-	52	0.6			
Belgium	-1.1	-3.5	62%	6.6%	23.3%	52%	6%	52	4.5%	-	-			
Denmark	3.9	-5.0	60%	9.0%	9.5%	61%	7%	9	2.1%	-	-			
Finland	-4.1	-7.4	57%	1.4%	52.9%	10%	7%	3	0.8%	38	0.7			
France	-5.3	-4.9	60%	5.6%	23.2%	45%	7%	-	-	428 <sup>d</sup>	0.7			
Germany	-5.9	-7.7	60%	3.8%	28.2%	44%	8%	-	-	350 <sup>d</sup>	0.4			
Greece	-2.9	2.3	64%	8.1%	28.5%	28%	10%	-	-	75 <sup>d</sup>	0.7			
Ireland	-3.2	0.5	60%	6.2%	19.8%	34%	10%	2	0.4%	15	0.4			
Italy	-1.8	-2.9	60%	3.9%	25.1%	44%	7%	-	-	-	-			
Luxembourg	-	-	57%	2.7%	10.8%	68%	0%	-	-	1	0.2			
Netherlands	-2.9	-4.1	59%	4.4%	16.9%	66%	4%	61	5.3%	34 <sup>d</sup>	0.2			
Portugal	-4.5	-5.0	68%	2.8%	15.1%	12%	10%	546	15.4%	211 <sup>e</sup>	2.0			
Spain	-1.0	-2.7	65%	6.4%	17.7%	19%	7%	394	5.0%	334 <sup>d</sup>	0.8			
Sweden	-3.9	4.2	53%	6.6%	21.8%	73%	5%	-	-	16 <sup>d</sup>	0.2			
United Kingdom	1.7	3.7	55%	5.5%	14.9%	65%	7%	548 <sup>e</sup>	8.3%	384 <sup>d</sup>	0.6			
<b>Subtotal 'EU-15'</b>	<b>-3.1%</b>	<b>-3.0%</b>	<b>61%</b>	<b>5.1%</b>	<b>20.6%</b>	<b>41%</b>	<b>7%</b>	<b>-</b>	<b>4.8%</b>	<b>-</b>	<b>0.6</b>			
<b>New EU countries (since 2004)</b>														
Bulgaria	2.2	0.8	67%	5.5%	19.5%	0%	6%	10 <sup>d</sup>	0.3%	268 <sup>d</sup>	3.4			
Cyprus	-	-	68%	8.1%	16.2%	68%	8%	-	-	-	-			
Czech Republic	-4.5	-6.8	64%	0.6%	37.3%	13%	3%	5 <sup>e</sup>	0.4%	68	0.7			
Estonia	6.7	-7.5	66%	0.2%	13.1%	16%	18%	33	6.4%	49	3.6			
Hungary	-3.4	-11.1	69%	0.3%	25.2%	3%	18%	-	-	191	1.9			
Latvia	7.4	-6.2	70%	4.7%	9.8%	6%	14%	51	3.5%	170	7.4			
Lithuania	5.7	-2.3	69%	3.5%	14.6%	3%	18%	7	0.3%	308 <sup>d</sup>	9.0			
Malta	-	-	88%	0.0%	24.0%	68%	4%	0	0.0%	1	0.2			
Poland	-6.3	-4.0	66%	1.1%	27.3%	0%	11%	15 <sup>e</sup>	0.1%	806	2.1			
Romania	4.1	1.9	70%	3.8%	10.8%	0%	24%	187	0.6%	2,089 <sup>d</sup>	9.6			
Slovakia	-6.3	-6.5	61%	2.9%	36.1%	4%	14%	1	0.1%	47	0.9			
Slovenia	-6.1	-5.9	58%	2.5%	33.8%	17%	10%	0	0.0%	17	0.6			
<b>Subtotal New EU countries</b>	<b>0.2%</b>	<b>-1.4%</b>	<b>68%</b>	<b>3.1%</b>	<b>16.2%</b>	<b>1%</b>	<b>19%</b>	<b>-</b>	<b>0.3%</b>	<b>-</b>	<b>2.1</b>			
<b>Subtotal all EU ('EU-27')</b>	<b>-1.6%</b>	<b>-2.4%</b>	<b>65%</b>	<b>4.0%</b>	<b>18.2%</b>	<b>19%</b>	<b>14%</b>	<b>-</b>	<b>0.7%</b>	<b>-</b>	<b>0.7</b>			



following an increase in 1995-2000, while in the Czech Republic, Hungary and Slovakia the decline was faster in the latter period than in earlier years. Bulgaria and Romania had smaller increases in rates in 2000-2005 than in 1995-2000.

In 2005, two thirds of notifications were among males, and rates among them increased after childhood, reaching a plateau in adulthood with a subsequent slight increase in old age (Figure 2). In females, rates were lower than in males after childhood, with a bi-modal pattern peaking in early adulthood and in old age. Rates decreased progressively in all age groups over time (Figure 3). This largely reflected the trend in many countries, including France, Germany, the Netherlands and Portugal. However, while rates in cases aged 55 or older decreased sharply in all countries, differences were noted in the progression of rates in children and young adults between countries (data not shown). In Romania (not included in Figure 3), rates increased in all age groups until 2002, after which they declined. In this country, the proportion of previously treated cases increased progressively in 1995-2005 to reach 24% in 2005, by far the highest in all the EU and West.

The Baltic States experienced a substantial rise in notification rates in the 1990s after their independence from the Soviet Union, which has only abated in recent years. In Latvia and Lithuania, which accounted for most cases from the Baltic States, TB notification rates increased in under-5 year olds, but decreased in adults in recent years. In the United Kingdom, rates remained stable at low levels in children but increased progressively in all age groups between 15 and 54, with a doubling in rates in the 25-34 year olds over the period, during which time the proportion of foreign TB cases increased steadily to 79% in 2005. The contribution of TB in immigrants to overall notification varied greatly between countries. While only 1% of cases reported by the central European countries

in 2005 were not autochthonous (new EU countries excluding Cyprus and Malta; country range: 0-17%), in the other countries 42% of TB cases were of foreign origin (range: 12-82%). In 27 countries with data for 2005, two thirds of cases of foreign origin were equally distributed between Asia and Africa, 19% were from another country of the EU and West or the Balkans, and 9% from a former Soviet Union republic outside the EU. Half the foreign cases originated from only 11 countries, which included high-burden countries from the Indian Sub-continent and Sub-Saharan Africa as well as other populous countries within the WHO European Region itself.


Aggregated data on HIV sero-status of TB cases reported in 2003 to 2005 were available for 21 countries. The highest HIV prevalence among TB cases was reported by Portugal (15%) and Iceland (9%, 1 case), and was 2-8% in 8 countries and 0-1% in 11 countries. HIV prevalence among TB cases was reportedly stable in 2000-2005 in most countries, but increased markedly in Estonia (from 0.1% to 6.4%) and Latvia (from 0.7% to 3.5%).

Total TB mortality rate was 1/100,000 population or less in most countries, but was higher in Bulgaria, Hungary, Poland, and Portugal (2-3), and even higher in the Baltic States and Romania (4-10). In the 'EU15' and West, mortality increased progressively by age, being more than four times higher in persons over 64 years when compared to those aged 55 to 64 years (Figure 4). In contrast, in the central European Member States, rates increased sharply from childhood to early middle-age but then more smoothly into old age.

### The Balkans


In 2005, 27,582 TB cases were reported by the seven Balkan countries, 74% of these cases by Turkey alone. The overall TB

**FIGURE 3**  
Age-group specific TB notification rates, EU & West\*, 1996-2004



\* excluding cases without information on age (769) and cases from countries without full age distribution for one or more years: Andorra, Bulgaria, Cyprus, Greece, Ireland, Monaco, San Marino, Spain. Romania excluded as the age-specific rates are very different from the rest of the EU and West.

**FIGURE 4**  
TB mortality rates by age-group, EU & West and East,\* latest available data



\* excluding 438 TB deaths with age unknown and including latest data (2003-2005) for countries using ICD-10 coding: 'EU15' and West (N=1,895): Austria, Finland, France, Germany, Israel, Luxembourg, Netherlands, Norway, Portugal, Spain, Sweden, Switzerland, United Kingdom; New EU (N=3,739): Czech Rep, Estonia, Hungary, Latvia, Lithuania, Poland, Romania, Slovakia, Slovenia; East (N=52,252): Belarus, Kazakhstan, Kyrgyzstan, Moldova, Russian Federation, Ukraine, Uzbekistan (Source: WHO Mortality Database, WHOSIS, update October 2007)

notification rate in the countries was 29/100,000, with rates higher in Bosnia and Herzegovina (55) than in the other countries (17-32). Between 2000 and 2005, rates decreased by 0.5-6.9% yearly in all countries, except for Turkey where they increased by 1.3%. A decline was observed in Bosnia and Herzegovina following an increase in the late 1990s.

Males predominated (range: 54-67%) among notifications. Children represented 5% of reported cases in 2005 but reached 14% in the Former Yugoslav Republic of Macedonia (overall rate 32/100,000). Notification rates in males increased sharply from childhood to early adulthood, and then increased slowly into old age. In females, rates were lower than in males in adults and decreased after early adulthood to rise again smoothly in the elderly.

As for central EU Member States, only 1% of cases overall were of foreign origin (9% in Croatia) and two thirds of these were from another Balkan country. HIV sero-prevalence among TB cases was reported by four countries and was low (range: 0.0-0.3%).

TB mortality rates were moderate in Croatia and the Former Yugoslav Republic of Macedonia (2.5 and 3.8/100,000 population respectively).

### East

In 2005, the overall TB notification rate in the 12 former Soviet Union republics in the East reached 110/100,000 population. Rates were higher than average in Kazakhstan (205), Moldova (162), Georgia (144) and Kyrgyzstan (130). Over half of the 306,015 cases reported in this area were from the Russian Federation. Notification rates in the East increased on average by 3.9% yearly between 2000 and 2005, but this ranged widely between countries (-3.9% to +19.9%). Much of the overall increase is attributable to increasing inclusion of previously treated cases, the proportion of which increased from 12% to 25% over this period as the number of new cases remained stable (Figure 1). However, the mean annual increase in rates was 2.5 times lower in 2000-2005 than that observed between 1995 and 2000.

While there was a male predominance among notified cases this is lower in the central Asian republics than in the other countries (57% vs. 72% respectively). Children accounted for 4% of the cases overall, but reached 13% in Kyrgyzstan among previously untreated cases. Most cases aggregated in the ages 15-44 years with only 7% of cases being over 64 years. Rates in males increased from childhood to middle age by a factor of 10 and then decreased sharply in the older ages. Notification rates in childhood were similar between the sexes but much higher in adult males than in females, in whom they peaked in young adulthood and in middle-age.

Most Eastern countries do not report TB in foreign citizens. In Moldova and the Russian Federation, foreign citizens represented 1% of notified cases. Information on TB-HIV comorbidity was sparse. Six countries reported results of HIV testing in TB patients, in which 0.2-2.0% of notified TB cases were HIV positive in recent years.

Mortality data for TB were available for nine countries, of which four had low coverage or completeness. Total TB mortality rates in the other five countries varied between 10.4 and 25.4/100,000 in the latest available year. Mortality rates increased rapidly from childhood to peak at age group 35-54 years, and then decreased at old ages.

### Discussion

The reversal in the decline in TB notification rates observed in the early 1990s in western countries of the WHO European Region persisted for some years but, with some notable exceptions, most countries have experienced a steady decrease in newly-diagnosed TB cases in recent years. As notifications among nationals decreased or remained stable in nearly all countries, cases of foreign origin came to represent a larger proportion of all TB cases reported. Immigration from countries with high TB prevalence has been one of the most important recent developments concerning TB in much of the industrialised EU. Thus, the recent increase in total TB cases observed in Sweden and the United Kingdom reflect the incremental trend in foreign-born cases as rates largely stabilised among the indigenous population. Populations of foreign origin generally experience higher TB notification rates than nationals [8]. The bi-modal pattern in age-specific notification rates in the western countries reflects the superimposition of patterns from foreign (largely young adult) and autochthonous (mostly elderly) populations. These changes in the profile of TB patients in the western countries are likely to impact negatively on treatment outcome. Foreign patients are more likely to have drug resistance [9]. Treatment interruption is more common in immigrants [10]. Furthermore, the increasing age of TB patients has a negative impact on likelihood of treatment success [11]. In Finland, deaths among TB cases - half of whom are nowadays elderly - are higher than elsewhere [12].

TB surveillance data and trends in the East have to be interpreted carefully, as in several countries TB notification has been influenced differently by changes in TB control systems since the early 1990s. Stabilisation or increase in notification rates may thus reflect improved case detection or changes in case definitions rather than actual incidence. The wide range in the proportion of notified cases having had TB in the past reflects differences in patient recruitment and in the definition of a notifiable case. Much of the increment in case reports in the East in recent years was due to the increased inclusion of previously treated cases while the number of newly diagnosed cases levelled off. This explains the spike in the Russian Federation in 2003, which influenced overall rates in the East given the large share of cases coming from this country alone (Figure 1). Similarly, the Baltic States started including forms of previously treated cases in addition to relapses midway in the period under study partly explaining a peaking of total notification rates around this time. Despite these artifacts, the high notification rates in young adults in the East still indicate intense transmission in recent years.

Testing for HIV among TB patients and for TB among HIV-positive individuals is problematic and comment on trends of TB/HIV comorbidity are impeded by incomplete information. In most countries the HIV status is known for only a small proportion of TB cases. The proportion of TB cases infected with HIV when calculated by including all notified TB cases in the denominator gives a conservative estimate in countries where HIV testing is offered only to a selection of patients based on risk. Retrieval of testing results is also incomplete. Notwithstanding, a number of observations can be made. In Balkan countries, the low prevalence of HIV among TB cases is associated with a low HIV prevalence in the general population of these countries up to now [13,14]. The increase in HIV prevalence among TB cases observed elsewhere reflects separate processes of particular concern. In the United Kingdom, there has been increasing immigration from countries with high prevalence for both TB and HIV in recent years [15]. In

western countries injecting drug users (IDUs) usually predominate among HIV cases with TB, suggesting a higher risk for developing TB among this HIV transmission group [16]. The same appears to be the case in the former Soviet Union republics where a dramatic increase in newly diagnosed HIV infections has occurred since the mid-1990s, mostly among IDUs [17]. Being an IDU was a strong predictor of HIV infection in younger adults with TB in Kiev city (Ukraine) in 2004-2005, and TB/HIV comorbidity among IDUs has reportedly increased since 2002 [18]. In 2002-2003, 92% of 49 TB/HIV cases detected in a cross-sectional survey of St Petersburg (Russian Federation) had injected drugs [19]. In Estonia, which has endured a sharp HIV-epidemic in the early years of this decade mostly in IDUs, [20] the steady rise in TB/HIV comorbidity is likely to represent an overlap of the HIV and TB epidemics in the indigenous population. And in Latvia, where cases having TB and HIV increased, 31 of 51 TB/HIV cases detected in 1998-2001 were in IDUs [21].

TB mortality rates follow roughly the same geographic gradient as TB morbidity. All former Soviet Union republics had high TB death-to-notification ratios. In contrast to countries further west mortality rates and death-to-notification ratios peaked before old age. This may reflect a higher lethality due to drug-resistant disease, the prevalence of which is high in countries like Kazakhstan [22]. In Ukraine, a country which has been particularly affected by the HIV epidemic for a number of years [13], this may be the effect of comorbidity. Despite these observations, some limitations are noted when comparing data between countries. The practice of coding the cause of death varies between vital registration systems in different countries. For instance Lithuania attributes much more TB deaths to military disease than neighbouring Estonia and Latvia. Most countries in the East never use codes for death from the late effects of TB (i.e. ICD-9 137 or ICD-10 B90), in contrast to many Western countries, some of which - like Norway and Sweden - register more deaths in these categories than under the standard tuberculosis codes used in this article [5].

### Conclusion

Countries in the East, with their high TB morbidity and mortality, remain a priority for TB surveillance in the WHO European Region. Surveillance systems need to be reinforced and modernized, and data collection increasingly automated. Access of TB patients to health care facilities with reliable laboratory facilities to perform culture examination and HIV testing, and the reporting of these test results should become more widespread. Cases of foreign origin should be more widely reported. The male predominance among adult TB cases, albeit more or less ubiquitous in the world [23], should serve as an alert to investigate possible barriers to access to care for women, especially when sex-ratios differ markedly between neighbouring countries with similar epidemiological patterns.

Today's EU presents a wide spectrum of TB patterns. Notification rates in the Baltic States remain high even if in decline, and these countries are particularly concerned by TB/HIV comorbidity and drug resistance. Central European countries, several bordering the former Soviet Union, need to be vigilant regarding a possible re-emergence of TB. The low TB incidence in much of the EU and West are no reason for complacency. The elimination of TB (<1 TB case/1,000,000 population) is still a distant prospect for all countries. The measurement of progress towards elimination will necessitate sensitive indices of disease activity in sub-populations at increased risk of TB infection. These groups need to be better

profiled at the supranational level. Similar to the way geographical origin has been built into international reporting, the collection of additional variables that are amenable to adequate standardisation can be useful for targeting public health action. These could include indices of social deprivation, history of imprisonment, use of tobacco, use of alcohol, injecting drug use, contact with active TB and area of residence within the country.

Despite efforts throughout the lifetime of EuroTB to standardise the European case definition for TB case reporting, more work is required. The Regional network of national TB surveillance authorities has provided a useful forum for discussion and exchange of experiences in the past, including the revision of the European case definition in 2006. This resource should be developed in future to enhance the process of routine data collection, conduction of surveys and standardisation of methodology.

### Acknowledgments

The EuroTB network was jointly funded by the European Commission (DG-SANCO) and the Institut de veille sanitaire of France. The following correspondents, in place at the end of 2007, had the opportunity to comment on an advanced draft of this article: M Coll Armangué (Andorra), H Hafizi (Albania), V Pogosian (Armenia), JP Klein (Austria), F Agaev (Azerbaijan), G Gurevich (Belarus), M Wanlin (Belgium), Z Dizdarevic, M Duronjic (Bosnia and Herzegovina), V Milanov (Bulgaria), A Simunovic (Croatia), C Hadjianastassiou (Cyprus), J Wallenfels (Czech Republic), P Andersen (Denmark), V Hollo (Estonia), P Ruutu (Finland), D Antoine (France), A Salakaia (Georgia), W Haas, B Brodhun (Germany), G Spala (Greece), J Strausz (Hungary), T Blondal (Iceland), J O'Donnell (Ireland), D Chemtob (Israel), MG Pompa (Italy), Sh Ismailov (Kazakhstan), A Alisherov (Kyrgyzstan), J Leimans (Latvia), E Davidaviciene (Lithuania), P Huberty-Krau (Luxembourg), S Talevski (the Former Yugoslav Republic of Macedonia), A Pace Ascjak (Malta), D Sain (Moldova), A Negre (Monaco), O Bojovic (Montenegro), C Erkens (the Netherlands), B Winje-Askeland (Norway), M Korzeniewska-Koseła (Poland), A Fonseca Antunes (Portugal), D Chiotan (Romania), E Kakorina (Russian Federation), A Sorcinelli (San Marino), G Radosavljevic-Asic (Serbia), I Solovic (Slovakia), D Erzen (Slovenia), E Rodriguez Valin (Spain), V Romanus (Sweden), P Helbling (Switzerland), S Saidaliev (Tajikistan), F Gümüslü (Turkey), B Jumaev (Turkmenistan), M Golubchikov (Ukraine), J Watson, J McMenamin, R Salmon, B Smyth (United Kingdom), D Usmanova (Uzbekistan). Likewise other members of the EuroTB Advisory Committee not already mentioned: L Clancy, F Drobniowski, E Ibrahim, M Forssbohm, V Kuyvenhoven and R Zaleskis. We also acknowledge the comments on this article made by J-C Desenclos and D Che of the Department of Infectious Diseases at the Institut de veille sanitaire.

### References

1. Raviglione MC, Sudre P, Rieder HL, Spinaci S, Kochi A. Secular trends of tuberculosis in Western Europe. *Bull World Health Organ* 1993; 71(3-4):297-306.
2. Coker RJ, Atun RA, McKee M. Health-care system frailties and public health control of communicable disease on the European Union's new eastern border. *Lancet* 2004;363(9418):1389-92.
3. Raviglione M, Rieder HL, Styblo K, Khomenko AG, Esteves K, Kochi A. Tuberculosis trends in Eastern Europe and the former USSR. *Tuber Lung Dis.* 1994 Dec;75(6):400-16.
4. United Nations Population Division. *Annual Populations 1950-2050 (The 2006 Revision)*, United Nations, New York. 2007.
5. World Health Organization. *Mortality Database*. Available from: <http://www3.who.int/whosis/en> (updated 15 October 2007, accessed 18 December 2007).

6. Rieder HL, Watson JM, Raviglione MC, Forssbohm M, Migliori GB, Schwoebel V, et al. Surveillance of tuberculosis in Europe. Working Group of the World Health Organization (WHO) and the European Region of the International Union Against Tuberculosis and Lung Disease (IUATLD) for uniform reporting on tuberculosis cases. *Eur Respir J*. 1996;9(5):1097-104.
7. The Commission of the European Communities. Commission Decision of 19 March 2002 laying down case definitions for reporting communicable diseases to the Community network under Decision No 2119/98/EC of the European Parliament and of the Council (2002/253/EC). *Official Journal of the European Communities* 2002; 86:44-62.
8. Infuso A, Falzon D, on behalf of the EuroTB network. [European surveillance of tuberculosis: description of the network and recent results] *Med Mal Infect*. 2005;35(5):264-8. French.
9. Falzon D, Infuso A, Ait-Belghiti F, for the EuroTB correspondents. In the European Union, TB patients from former Soviet countries have a high risk of multidrug resistance. *Int J Tuberc Lung Dis*. 2006;10(9):954-8.
10. Borgdorff MW, Veen J, Kalisvaart NA, Broekmans JF, Nagelkerke N. Defaulting from tuberculosis treatment in the Netherlands: rates, risk factors and trend in the period 1993-1997. *Eur Respir J*. 2000;16(2):209-13.
11. Falzon D, Le Strat Y, Belghiti F, Infuso A, for the EuroTB correspondents. Exploring the determinants of treatment success for tuberculosis cases in Europe. *Int J Tuberc Lung Dis* 2005;9(11):1224-1229.
12. Vasankari T, Holmström P, Öllgren J, Lippo K, Kokki M, Ruutu P. Risk factors for poor tuberculosis treatment outcome in Finland: a cohort study. *BMC Public Health*. 2007;7(147):291.
13. EuroHIV. HIV/AIDS surveillance in Europe. End-year report 2005. No. 73 ([www.eurohiv.org](http://www.eurohiv.org)). Saint-Maurice: Institut de veille sanitaire; 2006. Available from: [http://www.eurohiv.org/reports/report\\_73/pdf/report\\_eurohiv\\_73.pdf](http://www.eurohiv.org/reports/report_73/pdf/report_eurohiv_73.pdf)
14. UNAIDS Annual Report 2006: making the money work. Geneva: UNAIDS; 2006. Available from: [http://data.unaids.org/pub/Report/2007/2006\\_unaids\\_annual\\_report\\_en.pdf](http://data.unaids.org/pub/Report/2007/2006_unaids_annual_report_en.pdf)
15. Ahmed AB, Abubakar I, Delpech V, Lipman M, Boccia D, Forde J, et al. The growing impact of HIV infection on the epidemiology of tuberculosis in England and Wales: 1999-2003. *Thorax*. 2007;62(8):672-6.
16. Kirk O, Gatell JM, Mocroft A, Pedersen C, Proenca R, Brettler RP, et al. Infections with *Mycobacterium tuberculosis* and *Mycobacterium avium* among HIV-infected patients after the introduction of highly active antiretroviral therapy. EuroSIDA Study Group JD. *Am J Respir Crit Care Med*. 2000;162(3 Pt 1):865-72.
17. Hamers FF, Downs AM. HIV in central and eastern Europe. *Lancet* 2003;361(9362):1035-1044.
18. van der Werf MJ, Yegorova OB, Chentsova N, Chechulin Y, Hasker E, Petrenko VI, et al. Tuberculosis-HIV co-infection in Kiev City, Ukraine. *Emerg Infect Dis*. 2006;12(5):766-8.
19. Van Rie A, Zhemkov V, Granskaya J, Steklova L, Shpakovskaya L, Wendelboe A, et al. TB and HIV in St Petersburg, Russia: a looming catastrophe? *Int J Tuberc Lung Dis*. 2005;9(7):740-5.
20. Uuskula A, Kalikova A, Zilmer K, Tammai L, DeHovitz J. The role of injection drug use in the emergence of human immunodeficiency virus infection in Estonia. *Int J Infect Dis* 2002;6(1):23-27.
21. Morozova I, Riekstina V, Sture G, Wells C, Leimane V. Impact of the growing HIV-1 epidemic on multidrug-resistant tuberculosis control in Latvia. *Int J Tuberc Lung Dis* 2003;7(9):903-906.
22. World Health Organization/International Union Against TB and Lung Disease Global Project on Anti-Tuberculosis Drug Resistance Surveillance. Anti-Tuberculosis Drug Resistance in the World: Fourth Global Report. 2008. Available from: [http://www.who.int/tb/publications/2008/drs\\_report4\\_26feb08.pdf](http://www.who.int/tb/publications/2008/drs_report4_26feb08.pdf)
23. Holmes CB, Hausler H, Nunn P. A review of sex differences in the epidemiology of tuberculosis. *Int J Tuberc Lung Dis* 1998;2(2):96-104.

This article was published on 18 March 2008.

Citation style for this article: Falzon D, van Cauteren D. Demographic features and trends in tuberculosis cases in the European Region, 1995-2005. *Euro Surveill*. 2008;13(12):pii=8075. Available online: <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=8075>