Community-wide outbreaks of haemolytic uraemic syndrome associated with Shiga toxin-producing *Escherichia coli* O26 in Italy and Romania: a new challenge for the European Union

E Severi¹, F Vial¹, E Peron^{2,3}, O Mardh¹, T Niskanen¹, J Takkinen¹

1. European Centre for Disease Prevention and Control (ECDC), Stockholm, Sweden

2. European Programme for Intervention Epidemiology Training (EPIET), European Centre for Disease Prevention and Control (ECDC), Stockholm, Sweden

3. Gastrointestinal, zoonosis and tropical diseases unit, Department of infectious diseases epidemiology, Robert Koch Institute, Berlin, Germany

Correspondence: Ettore Severi (ettore.severi@ecdc.europa.eu)

Citation style for this article:

Severi E, Vial F, Peron E, Mardh O, Niskanen T, Takkinen J. Community-wide outbreaks of haemolytic uraemic syndrome associated with Shiga toxin-producing *Escherichia coli* O26 in Italy and Romania: a new challenge for the European Union. *Euro Surveill.* 2016;21(49):pii=30420. DOI: <http://dx.doi.org/10.2807/1560-7917.ES.2016.21.49.30420>

Article submitted on 02 December 2016 / accepted on 08 December 2016 / published on 08 December 2016

To the editor: In their recent article in *Eurosurveillance*, Germinario et al. describe a community-wide outbreak of Shiga toxin 2-producing *Escherichia coli* (STEC) O26:H11 infections associated with haemolytic uraemic syndrome (HUS) and involving 20 children between 11 and 78 months of age in southern Italy during the summer 2013 [1]. The investigation identified an association between STEC infection and consumption of dairy products from two local milk-processing establishments. We underline striking similarities to a recent multi-country STEC O26 outbreak in Romania and Italy and discuss the challenges that STEC infections and their surveillance pose at the European level.

In March 2016, Peron et al. published, also in *Eurosurveillance*, early findings of the investigation of a community-wide STEC infection outbreak in southern Romania [2]. As at 29 February 2016, 15 HUS cases with onset of symptoms after 24 January 2016, all but one in children less than two years of age, had been identified, three of whom had died. Aetiological confirmation was retrospectively performed through serological diagnosis and six cases were confirmed with STEC O26 infection. Shortly after this publication, and following the identification of the first epidemiologically-linked case in central Italy, the European Centre for Disease Prevention and Control (ECDC) and the European Food Safety Authority (EFSA) published a joint Rapid Outbreak Assessment [3]. The Italian and Romanian epidemiological, microbiological and environmental investigations implicated products from a milk-processing establishment in southern Romania as a possible source of infection. The dairy plant exported milk products to at least four European Union (EU)

countries. The plant was closed in March 2016 and the implicated food products recalled or withdrawn from the retail market.

Pulsed Field Gel Electrophoresis (PFGE) and whole genome sequencing (WGS) analyses did not establish a microbiological link between the Italian (2013) and the Romanian/Italian (2016) outbreaks (personal communication, Stefano Morabito, October 2016). However, the epidemiological similarities between the two community-wide outbreaks associated with HUS and STEC O26 infections, mostly affecting young children and implicating dairy products, are notable. While raw milk and unpasteurised dairy products are well known potential sources of STEC infection, milk products, as highlighted by Germinario et al. [1], have been rarely implicated in community-wide STEC outbreaks in the past, emphasising an emerging risk of STEC O26 infection associated with milk products.

Reporting of STEC O26 infections has been steadily increasing in the EU since 2007, partly due to improved diagnostics of non-O157 sero-pathotypes [4]. The attention to non-O157 STEC sero-pathotypes rose considerably after the severe STEC O104 outbreak that took place in Germany and France in 2011 during which almost 4,000 cases and more than 50 deaths were reported [5]. In light of the recently published outbreaks related to dairy products and the simultaneous increased reporting of isolations of STEC O26 from milk and milk products in the EU/European Economic Area (EEA) [6], strengthening STEC surveillance in humans and food and enhancing HUS surveillance in children less than five years of age is warranted. Paediatric

nephrologists should be sensitised to this effect and further joint studies between food and public health sectors be increased.

Conflict of interest

None declared.

Authors' contributions

ES drafted the letter to the Editor. All authors reviewed, commented and accepted its final version.

References

1. Germinario C, Caprioli A, Giordano M, Chironna M, Gallone MS, Tafuri S, et al. , all participants of the Outbreak investigation team. Community-wide outbreak of haemolytic uraemic syndrome associated with Shiga toxin 2-producing *Escherichia coli* O26:H11 in southern Italy, summer 2013. *Euro Surveill.* 2016;21(38):30343. DOI: 10.2807/1560-7917.ES.2016.21.38.30343 PMID: 27684204
2. Peron E, Zaharia A, Zota LC, Severi E, Mårdh O, Usein C, et al. Early findings in outbreak of haemolytic uraemic syndrome among young children caused by Shiga toxin-producing *Escherichia coli*, Romania, January to February 2016. *Euro Surveill.* 2016;21(11):30170. DOI: 10.2807/1560-7917.ES.2016.21.11.30170 PMID: 27020906
3. European Centre for Disease Prevention and Control (ECDC) and European Food Safety Authority. (EFSA). Multi-country outbreak of Shiga toxin-producing *Escherichia coli* infection associated with haemolytic uraemic syndrome – 5 April 2016. Stockholm: ECDC, 2016. Available from: <http://ecdc.europa.eu/en/publications/Publications/RRA-Escherichia-coli-O26-Romania-Italy-April2016.pdf>
4. European Centre for Disease Prevention and Control (ECDC). ECDC surveillance atlas of infectious diseases. Stockholm: ECDC. [Accessed 9 Oct 2016]. Available from: <http://ecdc.europa.eu/en/data-tools/atlas/pages/atlas.aspx>.
5. Frank C, Werber D, Cramer JP, Askar M, Faber M, an der Heiden M, et al. , HUS Investigation Team. Epidemic profile of Shiga-toxin-producing *Escherichia coli* O104:H4 outbreak in Germany. *N Engl J Med.* 2011;365(19):1771-80. DOI: 10.1056/NEJMoa1106483 PMID: 21696328
6. European Food Safety Authority and European Centre for Disease Prevention and Control. The European Union Summary Report on Trends and Sources of Zoonoses, Zoonotic Agents and Food-borne Outbreaks in 2012. *EFSA J.* 2014;12(2):3547. DOI: 10.2903/j.efsa.2014.3547

License and copyright

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0) Licence. You may share and adapt the material, but must give appropriate credit to the source, provide a link to the licence, and indicate if changes were made.

This article is copyright of the authors, 2016.