Indistinguishable NDM-producing *Escherichia coli* isolated from recreational waters, sewage, and a clinical specimen in Ireland, 2016 to 2017

BM Mahon¹, C Brehony¹, E McGrath^{2,3}, J Killeen¹, M Cormican^{1,2,3}, P Hickey⁴, S Keane⁴, B Hanahoe³, A Dolan⁵, D Morris¹

1. Antimicrobial Resistance and Microbial Ecology Group, School of Medicine, National University of Ireland, Galway, Ireland

2. Carbapenemase-Producing Enterobacteriaceae Reference Laboratory, Department of Medical Microbiology, University Hospital Galway, Galway, Ireland

3. Department of Medical Microbiology, University Hospital Galway, Galway, Ireland

4. Environmental Health Service, HSE West, Galway, Ireland

5. Galway County Council, Galway, Ireland

Correspondence: Bláthnaid Mahon (b.mahon1@nuigalway.ie)

Citation style for this article:

Mahon BM, Brehony C, McGrath E, Killeen J, Cormican M, Hickey P, Keane S, Hanahoe B, Dolan A, Morris D. Indistinguishable NDM-producing *Escherichia coli* isolated from recreational waters, sewage, and a clinical specimen in Ireland, 2016 to 2017. *Euro Surveill.* 2017;22(15):pii=30513. DOI: <http://dx.doi.org/10.2807/1560-7917.ES.2017.22.15.30513>

Article submitted on 15 March 2017 / accepted on 12 April 2017 / published on 13 April 2017

In this study, New Delhi metallo-beta-lactamase (NDM)-producing *Enterobacteriaceae* were identified in Irish recreational waters and sewage. Indistinguishable NDM-producing *Escherichia coli* by pulsed-field gel electrophoresis were isolated from sewage, a fresh water stream and a human source. NDM-producing *Klebsiella pneumoniae* isolated from sewage and seawater in the same area were closely related to each other and to a human isolate. This raises concerns regarding the potential for sewage discharges to contribute to the spread of carbapenemase-producing *Enterobacteriaceae*.

We report the finding of New Delhi metallo-beta-lactamase (NDM)-producing *Enterobacteriaceae* in fresh water and seawater samples collected at two beaches located near an untreated human sewage ocean discharge. Isolates of NDM-producing *Escherichia coli* derived from the sewage collection system, the sewage storage tank and the outflow were 100% identical by pulsed-field gel electrophoresis (PFGE) to those derived from a fresh water stream on one of the beaches, and to a clinical isolate.

Recreational water and sewage sample sites

In 2016, we identified a beach (Beach A) in Ireland, used for bathing and recreation, which is crossed by two fresh water streams (Stream A and Stream B), flowing from the surrounding countryside. These streams were examined for the presence of carbapenemase-producing *Enterobacteriaceae* (CPE). The detection of NDM-producing *E. coli* in these waters prompted subsequent additional sampling of the streams. As untreated human sewage was being discharged into the sea in the vicinity of the beach, and the fresh water streams

can become immersed in seawater at high tide, sewage was evaluated as a potential source. Sewage samples included samples from the collection system, the storage tank and the outflow. Sampling was performed in the period May to September, 2016. The sewage system is not linked to any hospital or long-term care facility that we are aware of. Further sampling of the fresh water streams and sewage sites was carried out in January 2017. In addition to this, seawater from Beach A and from a second beach (Beach B), ca 950 m in a direct line from Beach A were examined. Figure 1 shows a schematic diagram of the sampling points and their location relative to each other.

Processing of samples

We applied a previously described method (CapE), to examine large volumes of water (30L) from both the fresh water streams and the seawater, for the presence of CPE [1]. Following filtration and overnight enrichment, the samples were sub-cultured onto Brilliance CRE agar (Oxoid). Sewage samples were examined by direct plating onto Brilliance CRE agar. Following purification, presumptive isolates were identified to species level by matrix-assisted laser desorption/ionisation time-of-flight (MALDI-TOF) mass spectrometry, and antimicrobial susceptibility testing was performed and interpreted in accordance with European Committee on Antimicrobial Susceptibility Testing (EUCAST) criteria [2]. Carbapenemase-encoding genes were detected by real-time PCR, as previously described [3-6]. Typing of NDM-producing *Enterobacteriaceae* was not performed at the variant level, but PFGE was performed on all isolates, as outlined previously [7]. PFGE profiles of NDM-producing *Enterobacteriaceae* isolated from recreational water and sewage samples were compared with

FIGURE 1
Schematic diagram of water and sewage sampling points and their location relative to each other, Ireland, 2016–2017

FIGURE 2
PFGE analysis of New Delhi metallo-beta-lactamase-producing *Escherichia coli* isolated from fresh water, sewage and a clinical source in Ireland, 2016–2017 (n = 6 isolates)

PFGE: pulsed-field gel electrophoresis.

PFGE typing was performed with *Xba*1. All isolates are 100% similar by PFGE.

PFGE profiles of NDM-producing *Enterobacteriaceae* isolated from clinical specimens.

Findings of New Delhi metallo-beta-lactamase (NDM)-producing *Enterobacteriaceae*

Of eight fresh water samples from Stream B, NDM-producing *E. coli* were isolated from two samples, which were collected on 13 July and 24 August 2016. NDM-producing *E. coli* were also isolated in samples collected on 15 September 2016 from the sewage collection system (one of two samples), the storage tank (one of two samples) and the sewage outflow (one of one sample) (Table). All isolates were resistant to

ampicillin, cefotaxime, cefoxitin, cefpodoxime, cef-tazidime, ciprofloxacin, ertapenem, meropenem and nalidixic acid. The isolates obtained from the fresh water and sewage samples are indistinguishable by PFGE analysis from a human isolate submitted to the National Carbapenemase Producing *Enterobacteriaceae* Reference Laboratory Service (CPERLS) in early 2016 (Figure 2).

NDM-producing *K. pneumoniae* were isolated from two of three sewage sampling sites on 15 September 2016 and two of two sewage sampling sites on 18 January 2017. NDM-producing *K. pneumoniae* was also detected in seawater samples collected at Beach A and Beach B

FIGURE 3

PFGE analysis of New Delhi metallo-beta-lactamase-producing *Klebsiella pneumoniae* isolated from seawater, sewage and a clinical source in Ireland, 2016–2017 (n = 11 isolates)

PFGE: pulsed-field gel electrophoresis.

PFGE typing was performed with *Xba*I. Isolates are between 83% and 97% similar by PFGE

on 18 January 2017 (Table). These isolates were resistant to ampicillin, cefotaxime, ceftazidime, ciprofloxacin, ertapenem, gentamicin, kanamycin, meropenem, nalidixic acid and tetracycline. PFGE analyses of isolates from sewage, seawater and a human isolate (from CPERLS) show isolates to be between 83% and 97% similar (Figure 3).

Discussion

The rapid dissemination of carbapenemase-producing *Enterobacteriaceae* (CPE) in Europe and worldwide is making the delivery of effective healthcare an increasing challenge [8]. The number of CPE confirmed by the national reference laboratory in humans in Ireland has increased every year, rising from 48 in 2013, to 369 in 2016. In 2016, the three most commonly reported carbapenemases in Ireland were *K. pneumoniae* carbapenemase (KPC), carbapenem-hydrolysing oxacillinase-48 (OXA-48), and NDM [9]. The NDM gene is primarily plasmid-encoded, enabling its easy transfer between bacterial species. The plasmids are diverse and usually harbour a large number of other resistance genes [10]. The NDM gene has been detected extensively in the Indian subcontinent where it has been reported from both environmental and clinical sources [11]. Rapid global spread has been aided greatly by intercontinental travel [12]. However, a recent study in 2016 reported an outbreak of NDM-1-producing *Enterobacteriaceae* in a number of hospitals in Ireland, where links to foreign travel were not identified [13].

In Europe, a number of studies have reported the presence of CPE in recreational water, including Verona integron-encoded metallo-beta-lactamase (VIM) producing *K. pneumoniae* in a river in Switzerland in 2013 [14], KPC-producing *E. coli* in a river in Portugal in 2012 [15], VIM-1, VIM-34, and IMP-type metallo-beta-lactamase (IMP)-8 producing *E. coli* in the same Portuguese river in 2016 [16], and NDM-1 producing *K. pneumoniae* in the River Danube in Serbia, in 2016 [17]. Here we identify NDM-producing *Enterobacteriaceae* in environmental water samples collected at two adjacent beach sites in Ireland. As far as we are aware, this is the first such finding in bathing seawater in Europe.

We consider that contamination of the environment with NDM-producing *Enterobacteriaceae* from the human sewage outflow is likely to be the source, and that the fresh water streams were contaminated by backwash of sewage onto the beach by tidal currents. The presence of NDM-producing *Enterobacteriaceae* in the bathing water (seawater) and at a separate bathing site ca 950 m in a direct line indicates the extent of this contamination. It is important to note that by the established regulatory standards, the bathing water quality in the area concerned has been consistently of sufficient quality [18]. Notwithstanding compliance with regulatory standards however, it is reasonable to conclude that those using a beach such as this for recreational purposes might be at least intermittently

TABLE

Overview of sampling sites, dates and detection of carbapenemase-producing *Enterobacteriaceae* in a coastal region in Ireland, 2016–2017

Sample site	Date of sampling	Carbapenemase-producing <i>Enterobacteriaceae</i>
Fresh water Stream A	25 May 2016	Not detected
	22 Jun 2016	Not detected
	13 Jul 2016	Not detected
	10 Aug 2016	Not detected
	24 Aug 2016	Not detected
	7 Sep 2016	Not detected
	15 Sep 2016	Not detected
	18 Jan 2017	Not detected
Fresh water Stream B	25 May 2016	Not detected
	22 Jun 2016	Not detected
	13 Jul 2016	NDM-producing <i>E. coli</i>
	10 Aug 2016	Not detected
	24 Aug 2016	NDM-producing <i>E. coli</i>
	7 Sep 2016	Not detected
	15 Sep 2016	Not detected
	18 Jan 2017	Not detected
Sewage storage tank	15 Sep 2016	NDM-producing <i>E. coli</i>
		NDM-producing <i>K. pneumoniae</i>
	18 Jan 2017	NDM-producing <i>K. pneumoniae</i>
Sewage collection system	15 Sep 2016	NDM-producing <i>E. coli</i>
		NDM-producing <i>K. pneumoniae</i>
	18 Jan 2017	NDM-producing <i>K. pneumoniae</i>
Sewage outflow	15 Sep 2016	NDM-producing <i>E. coli</i>
Seawater Beach A	18 Jan 2017	NDM-producing <i>K. pneumoniae</i>
Seawater Beach B	18 Jan 2017	NDM-producing <i>K. pneumoniae</i>

E. coli: *Escherichia coli*; *K. pneumoniae*: *Klebsiella pneumoniae*; NDM: New Delhi metallo-beta-lactamase.

exposed to NDM-producing *Enterobacteriaceae*. Although, to date, there is no evidence that NDM-producing *Enterobacteriaceae* has been acquired as a result of exposure to this beach environment, Leonard et al. have recently reported on the level of risk of exposure to antibiotic resistant bacteria in coastal waters and its relationship to different types of water sports [19].

It appears therefore that there is potential for environmental contamination to contribute to a transition of CPE from largely healthcare-associated organisms, to organisms affecting the general population and the veterinary sector. From a public health perspective, the findings focus attention on the need to accelerate programmes to cease discharge of untreated sewage into the environment. This practice should be unacceptable in the context of discharges in the vicinity of popular bathing and recreation areas where human exposure is highly likely.

We consider that our findings point to potential limitations of the use of *E. coli* as an indicator bacteria for bathing water quality based on the number of colony

forming units (CFU) per 100 mL [20]. In our view, this approach does not reflect the pathogenicity of some variants of *E. coli*, such as Shiga-toxigenic *E. coli* for which the infectious dose is very low, (<10 CFU/mL) [21].

Conflict of interest

None declared.

Authors' contributions

Sampling was carried out by Bláthnaid Mahon, Carina Brehony, Dearbháile Morris, Paul Hickey and Shane Keane. Ann Dolan facilitated gaining access to the sewage supply. Water samples were filtered by Carina Brehony, James Killeen and Bláthnaid Mahon. Screening of samples for CPE was carried out by Bláthnaid Mahon. MALDI-TOF, antimicrobial susceptibility testing and PFGE was completed out by Bláthnaid Mahon, and real-time PCR by Elaine McGrath. Elaine McGrath and Belinda Hanahoe provided access to the clinical isolates. Bláthnaid Mahon, Dearbháile Morris and Martin Cormican were responsible for the study design and coordination, and wrote the manuscript. All authors were involved in reviewing the manuscript and approved the final version.

References

1. Morris D, Kavanagh S, Carney K, MacDomhnaill B, Cormican M. CapE (capture, amplify, extract): A rapid method for detection of low level contamination of water with Verocytotoxinogenic *Escherichia coli* (VTEC). *Sci Total Environ*. 2016;563-564:267-72. DOI: 10.1016/j.scitotenv.2016.04.075 PMID: 27135590
2. The European Committee on Antimicrobial Susceptibility Testing (EUCAST). Breakpoint tables for interpretation of MICs and zone diameters. Växjö: EUCAST. Version 7.0; 2017. Available from: <http://www.eucast.org>
3. Swayne RL, Ludlam HA, Shet VG, Woodford N, Curran MD. Real-time TaqMan PCR for rapid detection of genes encoding five types of non-metallo- (class A and D) carbapenemases in Enterobacteriaceae. *Int J Antimicrob Agents*. 2011;38(1):35-8. DOI: 10.1016/j.ijantimicag.2011.03.010 PMID: 21549572
4. Centers for Disease Control and Prevention (CDC). Multiplex real-time PCR detection of *klebsiella pneumoniae* carbapenemase (KPC) and new delhi metallo- β -lactamase (NDM-1) genes. Atlanta: CDC. 2011;6-7.
5. Birkett CI, Ludlam HA, Woodford N, Brown DFJ, Brown NM, Roberts MTM, et al. Real-time TaqMan PCR for rapid detection and typing of genes encoding CTX-M extended-spectrum β -lactamases. *J Med Microbiol*. 2007;56(Pt 1):52-5. DOI: 10.1099/jmm.0.46909-0 PMID: 17172517
6. Geyer CN, Reisbig MD, Hanson ND. Development of a TaqMan multiplex PCR assay for detection of plasmid-mediated ampC β -lactamase genes. *J Clin Microbiol*. 2012;50(11):3722-5. DOI: 10.1128/JCM.02038-12 PMID: 22895035
7. Swaminathan B, Barrett TJ, Hunter SB, Tauxe RV, CDC PulseNet Task Force. PulseNet: the molecular subtyping network for foodborne bacterial disease surveillance, United States. *Emerg Infect Dis*. 2001;7(3):382-9. DOI: 10.3201/eid0703.017303 PMID: 11384513
8. Albiger B, Glasner C, Struelens MJ, Grundmann H, Monnet DL, European Survey of Carbapenemase-Producing Enterobacteriaceae (EuSCAPE) working group. Carbapenemase-producing Enterobacteriaceae in Europe: assessment by national experts from 38 countries, May 2015. *Euro Surveill*. 2015;20(45):30062. DOI: 10.2807/1560-7917.ES.2015.20.45.30062 PMID: 26675038
9. McGrath E, Cormican M. CPEaRLS ANNUAL REPORT 2015, National Carbapenemase Producing Enterobacteriaceae (CPE) Reference Laboratory Service; 2015.1-26. Available from: https://www.researchgate.net/publication/307601382_Annual_Report_2015_Carbapenemase_Producing_Enterobacteriaceae_Reference_Laboratory_of_Ireland
10. Nordmann P, Naas T, Poirel L. Global spread of carbapenemase-producing Enterobacteriaceae. *Emerg Infect Dis*. 2011;17(10):1791-8. DOI: 10.3201/eid1710.110655 PMID: 22000347
11. Bushnell G, Mitrani-Gold F, Mundy LM. Emergence of New Delhi metallo- β -lactamase type 1-producing enterobacteriaceae and non-enterobacteriaceae: global case detection and bacterial surveillance. *Int J Infect Dis*. 2013;17(5):e325-33. DOI: 10.1016/j.ijid.2012.11.025 PMID: 23332300
12. Berrazeg M, Diene S, Medjahed L, Parola P, Drissi M, Raoult D, et al. New Delhi Metallo-beta-lactamase around the world: an eReview using Google Maps. *Euro Surveill*. 2014;19(20):20809. DOI: 10.2807/1560-7917.ES2014.19.20.20809 PMID: 24871756
13. O'Connor C, Cormican M, Boo TW, McGrath E, Slevin B, O'Gorman A, et al. An Irish outbreak of New Delhi metallo- β -lactamase (NDM)-1 carbapenemase-producing Enterobacteriaceae: increasing but unrecognized prevalence. *J Hosp Infect*. 2016;94(4):351-7. DOI: 10.1016/j.jhin.2016.08.005 PMID: 27624807
14. Zurfluh K, Hächler H, Nüesch-Inderbinen M, Stephan R. Characteristics of extended-spectrum β -lactamase- and carbapenemase-producing Enterobacteriaceae Isolates from rivers and lakes in Switzerland. *Appl Environ Microbiol*. 2013;79(9):3021-6. DOI: 10.1128/AEM.00054-13 PMID: 23455339
15. Poirel L, Barbosa-Vasconcelos A, Simões RR, Da Costa PM, Liu W, Nordmann P. Environmental KPC-producing *Escherichia coli* isolates in Portugal. *Antimicrob Agents Chemother*. 2012;56(3):1662-3. DOI: 10.1128/AAC.05850-11 PMID: 22203588
16. Kieffer N, Poirel L, Bessa LJ, Barbosa-Vasconcelos A, da Costa PM, Nordmann P. VIM-1, VIM-34, and IMP-8 Carbapenemase-Producing *Escherichia coli* Strains Recovered from a Portuguese River. *Antimicrob Agents Chemother*. 2016;60(4):2585-6. DOI: 10.1128/AAC.02632-15 PMID: 26810648
17. Kittinger C, Lipp M, Folli B, Kirschner A, Baumert R, Galler H, et al. Enterobacteriaceae isolated from the River Danube: Antibiotic resistances, with a focus on the presence of ESBL and carbapenemases. *PLoS One*. 2016;11(11):e0165820. DOI: 10.1371/journal.pone.0165820 PMID: 27812159
18. Environmental Protection Agency (EPA). Bathing water quality in Ireland 2015. EPA; 2016. Available from: https://www.epa.ie/pubs/reports/water/bathing/BW_Report_2015.pdf
19. Leonard AFC, Zhang L, Balfour AJ, Garside R, Gaze WH. Human recreational exposure to antibiotic resistant bacteria in coastal bathing waters. *Environ Int*. 2015;82:92-100. DOI: 10.1016/j.envint.2015.02.013 PMID: 25832996
20. Edberg SC, Rice EW, Karlin RJ, Allen MJ. *Escherichia coli*: the best biological drinking water indicator for public health protection. *Symp Ser Soc Appl Microbiol*. 2000;88(29):106S-16S. DOI: 10.1111/j.1365-2672.2000.tb05338.x PMID: 10880185
21. Croxen MA, Law RJ, Scholz R, Keeney KM, Wlodarska M, Finlay BB. Recent advances in understanding enteric pathogenic *Escherichia coli*. *Clin Microbiol Rev*. 2013;26(4):822-80. DOI: 10.1128/CMR.00022-13 PMID: 24092857

License and copyright

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0) Licence. You may share and adapt the material, but must give appropriate credit to the source, provide a link to the licence, and indicate if changes were made.

This article is copyright of the authors, 2017.