Universal screening for latent and active tuberculosis (TB) in asylum seeking children, Bochum and Hamburg, Germany, September 2015 to November 2016

Maya Mueller-Hermelink^{1,2}, Robin Kobbe^{2,3}, Benedikt Methling⁴, Cornelius Rau¹, Ulf Schulze-Sturm³, Isa Auer^{5,6}, Frank Ahrens¹, Folke Brinkmann⁴

1. Altona Children's Hospital, Department of Pulmonology, Hamburg, Germany

2. These authors contributed equally to this article

3. University Medical Centre Hamburg-Eppendorf, Department of Pediatrics, Hamburg, Germany

4. University Children's Hospital Bochum- Germany, Department of Pulmonology, Bochum, Germany

5. German Red Cross, Chapter Hamburg-Harburg, Hamburg, Germany

6. Center of Applied Sciences of Health (CASH), Leuphana University, Lueneburg, Germany

Correspondence: Folke Brinkmann (f.brinkmann@klinikum-bochum.de)

Citation style for this article:

Mueller-Hermelink Maya, Kobbe Robin, Methling Benedikt, Rau Cornelius, Schulze-Sturm Ulf, Auer Isa, Ahrens Frank, Brinkmann Folke. Universal screening for latent and active tuberculosis (TB) in asylum seeking children, Bochum and Hamburg, Germany, September 2015 to November 2016. *Euro Surveill*. 2018;23(12):pii=17-00536. <https://doi.org/10.2807/1560-7917.ES.2018.23.12.17-00536>

Article submitted on 31 Jul 2017 / accepted on 06 Mar 2018 / published on 22 Mar 2018

Background: In Germany, the incidence of tuberculosis (TB) in children has been on the rise since 2009. High numbers of foreign-born asylum seekers have contributed considerably to the disease burden. Therefore, effective screening strategies for latent TB infection (LTBI) and active TB in asylum seeking children are needed. **Aim:** Our aim was to investigate the prevalence of LTBI and active TB in asylum seeking children up to 15 years of age in two geographic regions in Germany. **Methods:** Screening for TB was performed in children in asylum seeker reception centres by tuberculin skin test (TST) or interferon gamma release assay (IGRA). Children with positive results were evaluated for active TB. Additionally, country of origin, sex, travel time, TB symptoms, TB contact and Bacille Calmette-Guérin (BCG) vaccination status were registered. **Results:** Of 968 screened children 66 (6.8%) had TB infection (58 LTBI, 8 active TB). LTBI prevalence was similar in children from high (Afghanistan) and low (Syria) incidence countries (8.7% vs 6.4%). There were no differences regarding sex, age or travel time between infected and non-infected children. Children under the age of 6 years were at higher risk of progression to active TB (19% vs 2% respectively, $p=0.07$). Most children (7/8) with active TB were asymptomatic at the time of diagnosis. None of the children had been knowingly exposed to TB. **Conclusions:** Asylum seeking children from high and low incidence countries are both at risk of developing LTBI or active TB. Universal TB screening for all asylum seeking children should be considered.

Introduction

In recent years, Germany experienced a major increase in the number of migrants. Of 890,000 refugees/

asylum seekers (hereafter referred to as asylum seekers) who reached Germany in 2015, more than one third were younger than 18 years. Most of them came from Syria, Iraq and Afghanistan [1].

Tuberculosis (TB) incidence rates in Germany had seen little change for several years until 2013, when numbers of TB cases started to rise. In 2015, an increase in incidence of 29.4% compared with the previous year was recorded [2]. Among the 5,865 cases notified there were 196 children up to 15 years of age. Incidence rates of TB are linked to migration with almost three quarters of TB patients in 2016 in Germany being foreign-born [3]. In foreign-born children the incidence of TB is 37 times higher than in children born in Germany (21.4 vs 0.6/100,000 children).

Asylum seekers have a higher risk of exposure to TB, both in their countries of origin and during migration. Furthermore, an increased risk for disease progression of latent tuberculosis infection (LTBI) due to physical and psychosocial stress factors has been observed [4]. In Germany, asylum seekers aged 15 years and older entering reception centres require a medical certificate stating the absence of any signs of potentially infectious pulmonary TB based on chest X-ray findings [5]. For children and adolescents up to the age of 15 years, an immunodiagnostic screening for TB, either by tuberculin skin test (TST) or interferon gamma release assay (IGRA), is recommended [6]. Immunodiagnostic tests can identify children with LTBI before they develop active TB [7]. In these children, preventive treatment significantly reduces the risk of progression and therefore further spread of the disease and further costs

FIGURE

Flowchart and results of screening procedures in children and adolescents at seven asylum seeker reception centres in Bochum and Hamburg, Germany, September 2015–November 2016 (n = 1,379)

IGRA: interferon gamma release assay; LTBI: latent tuberculosis infection; MMR: measles-mumps-rubella vaccination; TB: tuberculosis; TST: tuberculin skin test.

associated with severe TB, especially in young children [8,9].

The approach to screen all asylum seekers for LTBI and active TB would entail considerable pressure on healthcare resources. Responding to the European refugee crisis in 2015, many countries have therefore concentrated exclusively on screening for active TB [10] and so far, there is no uniform screening procedure for asylum seekers in Europe [10].

The aim of this study was to investigate the prevalence of TB infection in refugees up to 15 years of age in two geographic regions in Germany by immunological screening. The screening was carried out as part of the initial medical screening of asylum seekers at the reception centres ('Erstaufnahmeeinrichtung') upon arrival in Germany.

Methods

Between September 2015 and November 2016, children and adolescents aged from 3 months up to 15 years, were screened for TB infection using TST (2 IU PPD RT 23, Statens Serum Institute, Copenhagen, Denmark) and/or IGRA (QuantiFERON Gold in tube, Qiagen, Germantown, United States). Seven asylum seeker reception centres participated in two different German urban settings, three in Bochum and four in Hamburg. Asylum seekers in Germany usually stay in reception centres for 1 to 3 weeks before moving on to their secondary accommodation. Children who were immunised against measles, mumps and rubella (MMR) within the previous 4 weeks were excluded as live-attenuated MMR vaccines can lead to a temporary suppression of the cell-mediated immune response and result in false-negative TST results [11]. The excluded children were followed up 4 to 6 weeks later at the secondary accommodation.

A questionnaire assessed sociodemographic data including age, sex, country of origin, travel time as well as TB symptoms, TB contact (child of family member had contact with a known TB patient) and Bacille Calmette–Guérin (BCG) vaccination status.

In Hamburg, children were screened using TST and the test was considered positive if the transverse induration diameter was at least 10 mm as recommended by current guidelines [12]. In Bochum, children under the age of five were initially screened by TST, children and adolescents from 5 up to 15 years were screened by IGRA, due to a temporal shortage in TST. Screening and interpretation of TST results were the same as in Hamburg.

BCG vaccination status (mostly self-reported) and typical scars were recorded. Additionally, every child was assessed by a general medical examination. TST- or IGRA-positive children were referred to specialised children's hospitals for further diagnostics (e.g. chest X-ray, ultrasound, IGRA, sputum or gastric aspirates) and treatment.

Definitions

Following national guidelines, LTBI was defined as immunological evidence of infection (positive TST and/or IGRA) in absence of signs of active TB (clinical, radiological, microbiological) and active TB as evidence of TB with either microbiological and/or radiological and/or clinical signs of TB [6].

Statistical analysis

Data were analysed using SPSS 24.0 statistics software. Descriptive statistics were calculated for all variables. Comparisons between groups were made using the t-test or the Mann–Whitney U Test. For association between categorical variables chi-squared test or Fisher's exact test was performed. A p value less than 0.05 was considered to be statistically significant.

TABLE

Sociodemographic data of screened children by screening results in seven asylum seeker reception centres in Bochum and Hamburg, Germany, September 2015– November 2016

	Total		LTBI		Active TB		LTBI and active TB	
	N	%	n	%	n	%	n	%
Number of children	968	100	58	6	8	0.8	66	6.8
Age (median months)	71.1	NA	99	NA	55.5	NA	80.5	NA
Male	516	53.3	32	55.2	6	75	38	57.6
Country of origin								
Syria	377	38.9	21	36.2	3	37.5	24	36.4
Iraq	289	29.9	17	29.3	3	37.5	20	30.3
Afghanistan	217	22.4	17	29.3	2	25	19	28.8
Other ^a	85	8.8	3	5.2	0	0	3	4.5
BCG vaccinated	558	57.6	34	58.6	5	62.5	39	39.1
Travel time (median, in days)	21	NA	30	NA	30	NA	30	NA
TB contact	5	99.5	0	0	0	0	0	0
TB symptoms	69	7.1	2	3	1	12.5	1	0.02

BCG: Bacille Calmette-Guérin; LTBI: latent tuberculosis infection; NA: not applicable; TB: tuberculosis.

^a Other: Armenia (n= 1); Azerbaijan (n= 1); Georgia (n=5); Iran (n=41); Nigeria (n=11); Albania/Kosovo* and Serbia (n=14); Somalia (n=5); stateless (n=7).

*This designation is without prejudice to positions on status, and is in line with UNSCR 1244/1999 and the ICJ Opinion on the Kosovo declaration of independence.

Consent was provided by guardians in the presence of a translator. The study was approved by local ethical committees in both Bochum and Hamburg and conducted according to current guidelines for asylum seeking children and adolescent screening [12].

Results

A total of 1,379 asylum seeking children, aged from 3 months up to 15 years were included for evaluation in this study, 926 in Bochum and 453 in Hamburg.

We excluded 411 children from the screening: 217 had recently received MMR vaccine, 60 were screened previously in other reception centres, 80 had an acute febrile illness and 54 showed either indeterminate IGRA or were lost to follow-up (Figure). Recently vaccinated children, febrile children and those with indeterminate IGRA results received documentation regarding their current screening status and were (re-) screened at their long- term accommodation.

Of the 968 children who underwent screening the median age was 71.1 months (IQR 35–116). The majority (n=707; 73%) of the children were from the Middle East, mainly Syria (n=377; 38.9%) and Iraq (n=289; 29.9%). The third biggest group of asylum seeking children was from Afghanistan (n=217; 22.2%).

There were 66 children and adolescents with positive TST or IGRA results. Median age was 80.5 months (range: 6–168; IQR 42.7–141.5); 38 (57.6%) were boys. Of all infected children 24 (36.3%) were from Syria, 20

(30.3%) from Iraq and 19 (28.8%) from Afghanistan. In total, 31 children (47.7%) were under the age of 6 years at the time of screening. All 66 children and adolescents with positive TST or IGRA results were referred to a paediatric hospital for further diagnostics and therapy; 58 children (32 boys, 26 girls) were diagnosed with LTBI and treated with isoniazid and rifampicin for 3 months [13]. Eight children (6 boys, 2 girls) were diagnosed with active TB by chest X-ray and/or microbiological investigations and treated according to national recommendations [6]. Among these eight there were one sputum smear-positive adolescent and two children with PCR and microbiological cultures positive for *M. tuberculosis* complex.

Sociodemographic data and screening results are summed up in the Table.

Children with latent tuberculosis

Children with LTBI were older (median age 80.5 vs 71.1 months, $p=0.057$) than children without TB infection. The majority was from Syria (n=21; 36.2%), followed by Iraq (n=17; 29.3%) and Afghanistan (n=17; 29.3%). These percentages correlate well with the overall distribution of asylum seekers' countries of origin (Table).

The prevalence of LTBI differed only slightly between children from high and low incidence countries: Afghanistan (17/217; 7.8%) vs Syria (21/377; 5.6%; $p=0.27$) and Iraq (17/289; 5.9%; $p=0.44$). Although asylum seeking children with LTBI were older than the ones without LTBI, those under the age of 6 years were

at higher, although not statistically significant risk of progression to active TB (19% vs 2% respectively; $p=0,074$). There were no significant differences regarding sex, travel time, symptoms or BCG vaccination status between children with and without TB infection.

Children with active tuberculosis

The eight children with active TB were younger than the total of the screened children and those with LTBI (median 55.5 months vs 71.1 vs 80.5 months, $p=0,1$); six of them were male. Of the eight children, three were from Syria and three from Iraq. The other two were from Afghanistan. There were no significant differences in travel time and BCG vaccination in comparison to the total cohort. Seven children did not report any symptoms suggestive for TB (prolonged cough, fever, night sweats). None of them reported contact with a TB patient.

Discussion

Universal screening for TB detected 6.8 % children with LTBI and active TB in our study population of asylum seeking children up to the age of 15 years. About half of the children with LTBI were younger than 6 years and therefore at high risk of progression to active and especially disseminated TB if not treated promptly [14]. We also identified eight children in this age group who already progressed to active TB. In a large cohort study from Amsterdam, the risk of developing active TB was slightly lower for infected school age children but still reported to be up to 19.1% [15]. We identified two cases of active TB in this age group. Thus, our findings stress the importance of including children and young adolescents in national screening strategies.

Even though the overall risk of TB transmission by children is lower than by adults [14], there are reported cases in which school-aged children with active TB infected up to 39% of their contacts [16]. Considering the situation in crowded asylum seeker accommodations, the risk of transmission might be even higher [17]. The risk of developing active TB when infected has been shown to be higher in adult asylum seekers in comparison to the population in their countries of origin [18,19]. It remains unclear whether this is the case in children as well. An immunological screening of asylum seeking children of all age groups could potentially be helpful in preventing cases of infectious TB and in minimising the risk of progression to active and possibly disseminated disease in young children.

One approach to screening refugee children is to screen only children from countries with high TB incidence (above 100 cases/100,000 population). This approach does not seem ideal for different reasons [20]. Our data show that LTBI and active TB are not restricted to children from high incidence countries like Afghanistan (189 cases/100,000 population) [21]. Syrian children, the largest group of our study population, had similar percentages of TB infection as children from Afghanistan. The International Organization

for Migration (IOM) and the World Health Organization (WHO) documented rising TB incidence in Syria since 2013 [4]. For 2015, the official TB incidence is reported to be 20 cases per 100.000 inhabitants by the WHO, but it is doubtful whether this figure reflects the actual numbers as precise recording in times of war and civil unrest is known to be difficult [4].

Another reason for the increased numbers of TB infection in Syrian and Iraqi asylum seeking children might be related to travel routes and travel time to Germany. Of the 44 children from Syria and Iraq who had positive screening results, 13 spent a month or more in refugee camps along the Balkan route or before crossing the Mediterranean Sea. Very often medical support and nutrition during migration are inadequate and refugees from high and low TB incidence countries live together in crowded accommodations.

Symptom-based screening of refugees has been standard practice in many countries [10]. However, sensitivity and specificity of symptom-based screening is poor [22-24] and does not represent an effective screening approach, especially in young children [24]. In our study only one child with active TB displayed suggestive symptoms, while seven were detected by immunological screening and subsequent investigations.

Responding to the enormous increase in asylum seekers in 2015, many local authorities in Germany and other Western European countries concentrated on screening direct contacts of infectious TB patients [10,12]. Our data shows that contacts to TB cases were either not remembered or not stated by the children or their guardians, maybe due to fear of stigma connected with TB.

In our experience, universal TB screening in asylum seeking children is both effective and feasible. However, documentation of screening results and assurance of treatment completion in infected and diseased children remains crucial.

In addition to recording the screening result in the national vaccination card, asylum seeking children in the two urban areas involved in this study were supposed to receive a refugee health document (North Rhine-Westphalia health card (NRW Gesundheitskarte)); Hamburger health booklet (Hamburger Gesundheitsheft) in order to collate existing data and prevent redundant examinations. Despite the large overall amount of data that are collected in the process of providing healthcare to asylum seekers, there are neither federal standards for documentation, nor is there a standard set of collected health items in Germany [25,26]. In view of national and international mobility of asylum seekers, an innovative European solution of health information management would be desirable for the future.

Preventive therapy of LTBI in children is well tolerated and prevents progression to active TB in more than

90% [13]. Therefore, reduction of cases of active TB seems feasible in asylum seeking children. In order to assure treatment completion, different challenges have to be tackled in this hard-to-reach population. The stigma of being diagnosed with TB has to be addressed and resolved. Easy access to healthcare and increased tuberculosis awareness are vital [27]. In addition, directly observed therapy (DOTS) in asylum seeker reception centres and subsequent housing facilities could improve success rates.

Cost-effectiveness of TB screening was not investigated in our study. In a recent review in adults, LTBI screening in asylum seekers was described as cost effective [28]. None of the included studies, however, evaluated the subgroup of children.

An ideal approach would be to assess the overall exposure risk and perform baseline TB screening for all asylum seekers – both children and adults – upon arrival and pursue follow-ups over the next 2 years.

The ‘End TB Strategy’, adopted by WHO Member States in 2014, identifies TB in hard-to-reach populations – such as asylum seekers – as an important public health challenge for low-incidence countries like Germany and other Western European countries and recommends the implementation of comprehensive social and healthcare interventions [29].

Even though we screened a comparably high number of children, there are various study limitations that need to be taken into consideration. First our cohort is likely not to be representative for the general population of asylum seekers that reached Germany during the time of our investigation. It rather presents a snapshot at the moment of data collection, when the total numbers of asylum seekers, even in single reception centres, were not reliably documented. The distribution of asylum seekers within Germany was very heterogeneous for example regarding the country of origin and therefore a nationwide study would have been desirable. Another problematic issue was the high mobility of the asylum seekers. Anecdotal evidence showed that the latter were frequently registered twice at different centres and moved freely between them. Therefore, either double or incomplete registration could have caused a bias regarding epidemiological data, follow-up investigations and treatment [25,26].

Conclusion

In conclusion, our data supports the implementation of universal screening for LTBI and active TB in asylum seeking children of all ages and from both high and low TB incidence countries using immunological tests such as TST or IGRAs in order to prevent future active TB cases and further spread of the disease. Careful documentation of screening results and completion of preventive therapy should be ensured to guarantee the success of this screening approach.

Conflict of interest

The authors declare no conflict of interest.

Authors’ contributions

Maya Mueller-Hermelink, Robin Kobbe, Benedikt Methling, Cornelius Rau, Ulf Schulze-Sturm, Isa Auer, Frank Ahrens and Folke Brinkmann acquired clinical data. Maya Mueller-Hermelink, Robin Kobbe and Folke Brinkmann drafted the manuscript and Benedikt Methling, Cornelius Rau, Ulf Schulze-Sturm, Isa Auer, and Frank Ahrens reviewed the manuscript. All authors approved the final version.

References

1. Bundesamt für Migration und Flüchtlinge (BAMF). [Migration Report 2015 S 9-10. As of December 2016]. 2016. German. Available from: http://www.bamf.de/SharedDocs/Anlagen/DE/Publikationen/Migrationsberichte/migrationsbericht-2015.pdf?__blob=publicationFile
2. Brodhun B, Altmann D, Hauer B. [Report on the epidemiology of tuberculosis in Germany for 2015]. Berlin: Robert Koch Institut; 2016 German. Available from: https://www.rki.de/DE/Content/InfAZ/T/Tuberkulose/Download/TB2015.pdf?jsessionid=79CFDA71832AE18890A8273D33FAC506.1_cid390?__blob=publicationFile
3. Fiebig L, Hauer B, Brodhun B, Altmann D, Haas W. Tuberculosis in Germany: a declining trend coming to an end? *Eur Respir J*. 2016;47(2):667-70. <https://doi.org/10.1183/13993003.01410-2015> PMID: 26493803
4. Cookson ST, Abaza H, Clarke KR, Burton A, Sabrah NA, Ruman KA, et al. “Impact of and response to increased tuberculosis prevalence among Syrian refugees compared with Jordanian tuberculosis prevalence: case study of a tuberculosis public health strategy”. *Confl Health*. 2015;9(1):18. <https://doi.org/10.1186/s13031-015-0044-7> PMID: 26078784
5. [Law on the prevention and control of infectious diseases in humans]. (Infektionsschutzgesetz - IfSG), (2015). German. Available from: <https://www.gesetze-im-internet.de/ifsg/ifsg.pdf>
6. Schaberg T, Bauer T, Brinkmann F, Diel R, Feiterna-Sperling C, Haas W, et al. [Tuberculosis Guideline for Adults - Guideline for Diagnosis and Treatment of Tuberculosis including LTBI Testing and Treatment of the German Central Committee (DZK) and the German Respiratory Society (DGP)]. *Pneumologie*. 2017;71(6):325-97. PMID: 28651293
7. Heymann SJ, Brewer TF, Wilson ME, Colditz GA, Fineberg HV. Pediatric tuberculosis: what needs to be done to decrease morbidity and mortality. *Pediatrics*. 2000;106(1):E1. <https://doi.org/10.1542/peds.106.1.e1> PMID: 10878170
8. Morrison J, Pai M, Hopewell PC. Tuberculosis and latent tuberculosis infection in close contacts of people with pulmonary tuberculosis in low-income and middle-income countries: a systematic review and meta-analysis. *Lancet Infect Dis*. 2008;8(6):359-68. [https://doi.org/10.1016/S1473-3099\(08\)70071-9](https://doi.org/10.1016/S1473-3099(08)70071-9) PMID: 18450516
9. Rutherford ME, Hill PC, Triasih R, Sinfield R, van Crevel R, Graham SM. Preventive therapy in children exposed to *Mycobacterium tuberculosis*: problems and solutions. *Trop Med Int Health*. 2012;17(10):1264-73. <https://doi.org/10.1111/j.1365-3156.2012.03053.x> PMID: 22862994
10. Dara M, Solovic I, Sotgiu G, D'Ambrosio L, Centis R, Tran R, et al. Tuberculosis care among refugees arriving in Europe: a ERS/WHO Europe Region survey of current practices. *Eur Respir J*. 2016;48(3):808-17. <https://doi.org/10.1183/13993003.00840-2016> PMID: 27492827
11. Tamashiro VG, Perez HH, Griffin DE. Prospective study of the magnitude and duration of changes in tuberculin reactivity during uncomplicated and complicated measles. *Pediatr Infect Dis J*. 1987;6(5):451-4. <https://doi.org/10.1097/00006454-198705000-00007> PMID: 3601492
12. Ritz N, Brinkmann F. [Consensus based guideline “Tuberculosis Screening in asylum seeking children and teenagers under the age of 15” under direction of the German Society for Pediatric Infectiology (DGPI)]. *Monatsschrift für Kinderheilkunde*. 2015;163(12).
13. Spyridis NP, Spyridis PG, Gelesme A, Sypsa V, Valianatou M, Metsou F, et al. The effectiveness of a 9-month regimen of isoniazid alone versus 3- and 4-month regimens of isoniazid plus rifampin for treatment of latent tuberculosis infection in children: results of an 11-year randomized study. *Clin Infect*

- Dis. 2007;45(6):715-22. <https://doi.org/10.1086/520983> PMID: 17712755
14. Marais BJ, Gie RP, Schaaf HS, Hesselning AC, Obihara CC, Starke JJ, et al. The natural history of childhood intra-thoracic tuberculosis: a critical review of literature from the pre-chemotherapy era. *Int J Tuberc Lung Dis.* 2004;8(4):392-402. PMID: 15141729
 15. Sloot R, Schim van der Loeff MF, Kouw PM, Borgdorff MW. Risk of tuberculosis after recent exposure. A 10-year follow-up study of contacts in Amsterdam. *Am J Respir Crit Care Med.* 2014;190(9):1044-52. <https://doi.org/10.1164/rccm.201406-1159OC> PMID: 25265362
 16. Piccini P, Chiappini E, Tortoli E, de Martino M, Galli L. Clinical peculiarities of tuberculosis. *BMC Infect Dis.* 2014;14(Suppl 1):S4. <https://doi.org/10.1186/1471-2334-14-S1-S4> PMID: 24564419
 17. Beggs CB, Noakes CJ, Sleigh PA, Fletcher LA, Siddiqi K. The transmission of tuberculosis in confined spaces: an analytical review of alternative epidemiological models. *Int J Tuberc Lung Dis.* 2003;7(11):1015-26. PMID: 14598959
 18. Blount RJ, Tran MC, Everett CK, Cattamanchi A, Metcalfe JZ, Connor D, et al. Tuberculosis progression rates in U.S. Immigrants following screening with interferon-gamma release assays. *BMC Public Health.* 2016;16(1):875. <https://doi.org/10.1186/s12889-016-3519-6> PMID: 27558397
 19. Lönnroth K, Mor Z, Erkens C, Bruchfeld J, Nathavitharana RR, van der Werf MJ, et al. Tuberculosis in migrants in low-incidence countries: epidemiology and intervention entry points. *Int J Tuberc Lung Dis.* 2017;21(6):624-37. <https://doi.org/10.5588/ijtld.16.0845> PMID: 28482956
 20. Graham SM, Ahmed T, Amanullah F, Browning R, Cardenas V, Casenghi M, et al. Evaluation of tuberculosis diagnostics in children: 1. Proposed clinical case definitions for classification of intrathoracic tuberculosis disease. Consensus from an expert panel. *J Infect Dis.* 2012;205(Suppl 2):S199-208. <https://doi.org/10.1093/infdis/jis008> PMID: 22448023
 21. World Health Organization (WHO). Global tuberculosis report 2016. World Health Organization, Geneva; 2016. Available from: <http://apps.who.int/medicinedocs/documents/s23098en/s23098en.pdf>
 22. Cuevas LE, Petrucci R, Swaminathan S. Tuberculosis diagnostics for children in high-burden countries: what is available and what is needed. *Paediatr Int Child Health.* 2012;32(sup2) Suppl 2:S30-7. <https://doi.org/10.1179/2046904712Z.000000000076> PMID: 23394756
 23. Marais BJ, Gie RP, Hesselning AC, Schaaf HS, Enarson DA, Beyers N. Radiographic signs and symptoms in children treated for tuberculosis: possible implications for symptom-based screening in resource-limited settings. *Pediatr Infect Dis J.* 2006;25(3):237-40. <https://doi.org/10.1097/01.inf.0000202140.76368.74> PMID: 16511386
 24. Frigati L, Maskew M, Workman L, Munro J, Andronikou S, Nicol MP, et al. Clinical Predictors of Culture-confirmed Pulmonary Tuberculosis in Children in a High Tuberculosis and HIV Prevalence Area. *Pediatr Infect Dis J.* 2015;34(9):e206-10. <https://doi.org/10.1097/INF.0000000000000792> PMID: 26376315
 25. Bozorgmehr K, Nöst S, Thaiss HM, Razum O. [Health care provisions for asylum-seekers : A nationwide survey of public health authorities in Germany]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz.* 2016;59(5):545-55. <https://doi.org/10.1007/s00103-016-2329-4> PMID: 27072501
 26. Klinkhammer G, Korzilius H. Flüchtlinge: Gut versorgt- dank großer Hilfsbereitschaft. *German. Dtsch Arztebl.* 2015;41(112).
 27. de Vries SG, Cremers AL, Heuvelings CC, Greve PF, Visser BJ, Bèlard S, et al. Barriers and facilitators to the uptake of tuberculosis diagnostic and treatment services by hard-to-reach populations in countries of low and medium tuberculosis incidence: a systematic review of qualitative literature. *Lancet Infect Dis.* 2017;17(5):e128-43. [https://doi.org/10.1016/S1473-3099\(16\)30531-X](https://doi.org/10.1016/S1473-3099(16)30531-X) PMID: 28291721
 28. Zammarchi L, Casadei G, Strohmeyer M, Bartalesi F, Liendo C, Matteelli A, et al. . A scoping review of cost-effectiveness of screening and treatment for latent tuberculosis infection in migrants from high-incidence countries. *BMC Health Serv Res.* 2015;15(1):412. <https://doi.org/10.1186/s12913-015-1045-3> PMID: 26399233
 29. Heuvelings CC, de Vries SG, Greve PF, Visser BJ, Bèlard S, Janssen S, et al. Effectiveness of interventions for diagnosis and treatment of tuberculosis in hard-to-reach populations in countries of low and medium tuberculosis incidence: a systematic review. *Lancet Infect Dis.* 2017;17(5):e144-58. [https://doi.org/10.1016/S1473-3099\(16\)30532-1](https://doi.org/10.1016/S1473-3099(16)30532-1) PMID: 28291722

License and copyright

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0) Licence. You may share and adapt the material, but must give appropriate credit to the source, provide a link to the licence, and indicate if changes were made.

This article is copyright of the authors, 2018.