Sharing more than friendship – transmission of NDM-5 ST167 and CTX-M-9 ST69 *Escherichia coli* between dogs and humans in a family, Finland, 2015

Thomas Grönthal¹, Monica Österblad^{2,3}, Marjut Eklund¹, Jari Jalava³, Suvi Nykäsenoja⁴, Katariina Pekkanen⁴, Merja Rantala¹

1. Faculty of Veterinary Medicine, University of Helsinki, Helsinki, Finland

2. Sydspetsens miljöhälsa, Hangö, Finland

3. National Institute for Health and Welfare, Turku, Finland

4. Food Safety Authority Evira, Helsinki, Finland

Correspondence: Thomas Grönthal (thomas.gronthal@helsinki.fi)

Citation style for this article:

Grönthal Thomas, Österblad Monica, Eklund Marjut, Jalava Jari, Nykäsenoja Suvi, Pekkanen Katariina, Rantala Merja. Sharing more than friendship – transmission of NDM-5 ST167 and CTX-M-9 ST69 *Escherichia coli* between dogs and humans in a family, Finland, 2015. Euro Surveill. 2018;23(27):pii=1700497. <https://doi.org/10.2807/1560-7917.ES.2018.23.27.1700497>

Article submitted on 21 Jul 2017 / accepted on 03 Apr 2018 / published on 05 Jul 2018

Introduction: Carbapenemase-producing *Enterobacteria* (CPE) have rarely been reported in dogs, and never in animals in Finland. However, in April 2015, two meropenem-resistant *Escherichia coli* were identified from two dogs in one family. Both dogs suffered from chronic *otitis externa*. **Methods:** Epidemiological and molecular investigations (pulsed-field gel electrophoresis (PFGE), multilocus sequence typing) were conducted to investigate the source of infection and transmission routes. **Results:** In both dogs and one family member New Delhi metallo-beta-lactamase (NDM-5)-producing multidrug-resistant ST167 *E. coli* was found. Whole genome sequencing confirmed that the isolates were identical or only had one or two allelic differences. Additionally, the dogs and humans of the family carried an identical extended-spectrum beta-lactamase (ESBL) CTX-M-group 9 *E. coli* ST69 strain, indicating interspecies transmission. While the original source remains unclear, human-to-canine transmission is possible. No carbapenems had been administered to the dogs, but exposure to numerous other antimicrobials likely sustained the bacteria and supported its propagation in the canine host. **Conclusion:** To our knowledge, canine clinical NDM-5 *E. coli* in Europe, and confirmed CPE transmission between dogs and humans have not been previously reported. The screening of veterinary *Enterobacteriaceae* isolates for carbapenem resistance is highly recommended.

Introduction

The rapid global emergence of resistance to carbapenem and other extended-spectrum beta-lactams in *Enterobacteriaceae*, *Acinetobacter* and *Pseudomonas* species in the past decade is a major public health concern [1,2]. With such types of resistance, these species have the potential to cause severe infection, especially in hospitalised individuals [3]. Carbapenem resistance

is mediated by different types of carbapenemases, of which *Klebsiella pneumoniae* carbapenemase (KPC-type) and New Delhi metallo-beta-lactamase (NDM-type) carbapenemase are the most common in bacteria from human clinical specimens [2]. In addition to resistance to nearly all beta-lactams, isolates carrying carbapenemases are often multi- to extensively drug-resistant (MDR or XDR), leaving very few or no treatment options [4]. Carbapenemase-producing MDR bacteria were first described almost exclusively in humans, but since 2011 have also been detected in livestock, companion animals, wildlife, and different environmental compartments [5,6], indicating their transfer to new hosts and reservoirs. Carbapenem use in humans has been implicated as a cause of emerging carbapenem resistance [7]. Due to the MDR/XDR character of carbapenemase-producing bacteria, carbapenem resistance may also be co-selected for by the use of other antimicrobial classes [8]. Carbapenems are not authorised for veterinary use in Finland (Government decree 1054/2014) [9].

In Finland, carbapenemase-producing bacteria have been found in humans, although cases are still rare, with 20–34 new cases annually in 2013–16. Over half of these were probably of foreign origin [10]. However, such bacteria have never been identified in animals in the country. In the spring of 2015, meropenem-resistant *Escherichia coli* was diagnosed from two canine ear specimens in the Clinical Microbiology Laboratory (CML) of the Faculty of Veterinary Medicine, University of Helsinki. The exceptional findings initiated epidemiological and molecular investigations together with the Finnish Food Safety Authority (Evira) and the National Institute of Health and Welfare (THL). Here, we present the investigations and their results.

TABLE 1

Primers for extended-spectrum beta-lactamase, AmpC and carbapenemase multiplex PCRs and for sequencing of CTX-M group

PCRs	Forward (-F) and reverse (-R) primer sequences and orientation (5' → 3')	Target	Size (bp)	PCR conditions ^a	Reference
ESBL-multiplex	CTX-F: ATGTGCAGYACCAGTAARGTKATGGC	CTX-M	513	98°C (30s), 98°C (5s), 65°C (15s), 72°C (20s), 72°C (60s)	[19]
	CTX-R: CDCCGCTGCCGGTYTTATCVCC				[20]
	TEM-F: ATTTYCGTGTGCGCCTTATTCC	TEM	431		[20]
	TEM-R: AAGCGGTTAGCTCCTTCGGTC				This study
	SHV-F: ACCAGCCAGCGTCTGAGC	SHV	285		This study
	SHV-R: TTGCCAGTGCTCGATCAG				[21]
AmpC-multiplex	CIT-F: TGGCCAGAACTGACAGGCAAA	CIT	462	98°C (30s), 98°C (5s), 65°C (15s), 72°C (20s), 72°C (20s)	[22]
	CIT-R: TTTCTCCTGAACGTGGCTGGC				
	DHA-F: AACTTTCACAGGTGTGCTGGGT	DHA	405		
	DHA-R: CCGTACGCATACTGGCTTTGC				
	ACC-F: AACAGCCTCAGCAGCCGGTTA	ACC	345		
	ACC-R: TTCGCCGAATCATCCCTAGC				
	FOX-F: AACATGGGGTATCAGGGAGATG	FOX	190		
	FOX-R: CAAAGCGCGTAACCGGATTGG				
Carbapenemase-multiplex	KPC-for: CTTGCTGCCGCTGTGCTG	KPC	488	94°C (10 min), 94°C (30s), 59°C (30s), 72°C (60s)	[15]
	KPC-rev: GCAGGTTCCGGTTTTGTCTC				
	Imp-F-g-16: GCAGGTTCCGGTTTTGTCTC	IMP	232		
	Imp-R-g13–4: CCAAACyACTACGTTATCTkGAG				
	Vim-F: GATGGTGTGTTGGTCGCATA	VIM	389		
	Vim-R: CGAATGCGCAGCACCAG				
	NDM-for: GGCAGCACACTTCTATCTC	NDM	155		
	NDM-rev: GTTGATCTCCTGCTTGATCC				
	OXA-48A: TTGGTGGCATCGATTATCGG	OXA	743		
OXA-48B: GAGCACTTCTTTGTGATGGC					
	GES-1A: ATGCGCTTCATTCACGCAC	GES	863		
	GES-1B: CTATTTGTCCGTGCTCAGG				
	IMI-A: ATAGCCATCCTTGTTTAGCTC	IMI	818		
	IMI-B: TCTGCGATTACTTTATCCTC				
CTX-M gene grouping ^b	CTX-F: ATGTGCAGYACCAGTAARGTKATGGC	CTX-M	513	98°C (30s), 98°C (5s), 65°C (10s), 72 °C (15s), 72°C (60s)	[19]
	CTX-R: CDCCGCTGCCGGTYTTATCVCC				[20]

ESBL: extended-spectrum beta-lactamase.

^a Conditions for ESBL-, AmpC multiplex PCRs and for CTX-M-gene PCRs are reported in the following order: initial denaturation, denaturation, annealing, extension (30 cycles), and final extension.

^b The same primers were used for PCR and sequencing.

Methods

Epidemiological investigation

In spring 2015, two dogs, belonging to the same family, were found to have meropenem-resistant MDR *E. coli* in clinical specimens from their ears. Written consent for an epidemiological investigation was obtained from the owners. The investigation covered a timeline of up

to two years before the findings. The medical records of the dogs were obtained from the treating veterinarian. The owners were asked to fill in an electronic questionnaire that covered their medical history including hospitalisations, human and animal contacts, travel history and the living conditions of the dogs. The family members were also instructed to contact their local health-care centre to consider screening for carbapenem and

TABLE 2

Antimicrobial treatments^a the dogs had received before the NDM-*Escherichia coli* finding, Finland, 2013–2015

Dog A		Dog B	
Topical	Systemic	Topical	Systemic
Gentamicin	Amoxicillin-clavulanic acid	Fusidic acid	Cefalexin
Framycetin		Framycetin	Clindamycin
Fusidic acid	Cefalexin	Marbofloxacin	Enrofloxacin
Marbofloxacin	Enrofloxacin	Miconazole	Fluconazole
Miconazole	Itraconazole	Polymyxin B	Itraconazole
Polymyxin B			

NDM: New-Delhi metallo-beta-lactamase producing.

^a In addition, the dogs had received local and systemic corticosteroids.

extended-spectrum cephalosporin-resistant bacteria. The presence of carbapenemase and other extended-spectrum beta-lactamase-producing bacteria in the canine patients was followed for a year by taking specimens from the dogs' ears (4 occasions) and rectum (3 occasions). In addition, a retrospective search of the records of the CML and the Reference Laboratory of THL was performed for the presence of respective *E. coli* sequence types (ST) with similar antibiograms to those in this study.

Microbiological investigation

Laboratory methods for canine specimens

Clinical ear specimens were cultured on 5% sheep blood agar (tryptic soy agar with Sheep Blood, Oxoid, United Kingdom (UK)), a chromogenic agar (UriSelect 4, Bio-Rad, France) and Dixon agar plates (Tammertutkan Maljat, Finland). Bacteria were identified using biochemical methods [11], while disk diffusion susceptibility testing, including quality control and interpretation, was performed according to Clinical and Laboratory Standards Institute (CLSI) guidelines with *E. coli* ATCC 25922 as a quality control [12,13]. If no veterinary-specific susceptibility breakpoints were available, human CLSI breakpoints were used [14]. The susceptibility testing panel for *Enterobacteriaceae* included amikacin, amoxicillin-clavulanic acid, ampicillin, cefpodoxime, chloramphenicol, doxycycline, enrofloxacin, gentamicin, meropenem and trimethoprim-sulfamethoxazole (Oxoid, UK). Follow-up screening specimens from the ears and rectum were cultured on a selective agar (Brilliance ESBL, Oxoid, UK) and a blood agar plate with a 10 µg meropenem disk and incubated at $+35 \pm 1^\circ\text{C}$ for up to 48 hours. *Enterobacteriaceae* isolates growing on the selective plate or showing reduced susceptibility to cefpodoxime (≤ 22 mm) or meropenem (≤ 22 mm) were identified and preserved for further research. Preliminary screening of carbapenemase activity was carried out using the modified Hodge test [14]. The

type of beta-lactamase produced was confirmed by using commercial disk sets (carbapenemases: KPC+ MBL Confirm ID Kit, Rosco Diagnostica, Denmark; ESBL/AmpC-type beta-lactamases: AmpC and ESBL Detection Set, Mast Group Ltd, UK). Carbapenemase activity was confirmed with ultraviolet-spectrometric detection of imipenem hydrolysis [15].

Bacterial strains from humans

Three extended-spectrum beta-lactamase (ESBL)/carbapenemase-producing *E. coli* isolates isolated from rectal swabs of two human family members were received from the Helsinki-Uusimaa Hospital District Laboratory, HUSLAB. The rectal swabs were obtained and the specimens were cultured according to guidelines that have since been published [16]. In addition, susceptibility results were available from a fourth human ESBL *E. coli* isolated from a follow-up screening specimen. Sequence typing of the isolates was conducted as further described. A search from the National Infectious Diseases Register at THL was performed to identify human *E. coli* isolates with similar ST and carbapenemase profiles in Finland in 2014–15. These were included in the core genome multilocus sequence typing (cgMLST) comparison (see below).

Molecular methods

Pulsed-field gel electrophoresis

Pulsed-field gel electrophoresis (PFGE) was performed as previously described [17]. Clonal similarity was determined using the Dice coefficient and the unweighted pair-group method using arithmetic average clustering (UPGMA), with the optimisation and tolerance both set to 1% (GelCompar software version 6, Applied Maths, Belgium).

Multilocus sequence typing and detection of beta-lactamase-encoding genes by PCR

DNA for multilocus sequence typing (MLST) and PCR was extracted by suspending one to three bacterial colonies in 200 µL of PCR-grade water. The suspension was boiled for 10 min and centrifuged at 13,000 rpm for 5 min. The supernatant was used as a DNA template.

For MLST, amplification of *adk*, *fumC*, *gyrB*, *icd*, *mdh*, *purA* and *recA* genes was performed using primers by Lau et al. [18], which were attached to universal forward (UniF: 5'-GTTTCCCAGTCACGACGTTGTA-3') and reverse (UniR: 5'-TTGTGAGCGGATAACAATTTC-3') sequences at the 5' end of each primer. The universal F and R sequences were used as sequencing primers. The PCR reaction mixture (total volume 20 µL) contained 10 µL of 2x Phusion Flash High Fidelity Master Mix (Thermo Scientific, United States (US)), 0.25 µM of each primer (Oligomer Oy, Finland) and 1 µL of DNA template for all reactions, except for *icd*, for which a primer concentration of 0.5 µM was used. PCR conditions for *adk*, *fumC* and *gyrB* were as follows: initial denaturation at 98°C (15 s); 30 cycles of denaturation at 98°C (2 s), annealing at 63°C (10 s), elongation at

TABLE 3

Timeline and results of ear and screening specimens from dogs A and B, Finland, December 2013–May 2016 (n = 18 specimens)

Dog	Time	Specimen type	Finding	<i>E. coli</i> ID-number
A	March 2015	Ear	NDM- <i>Escherichia coli</i>	P-1009
	April 2015	Ear	NDM- <i>E. coli</i> , CoNS	P-1036
	June 2015	Ear	No growth	NA
	September 2015	Ear	<i>Moraxella</i> sp.	NA
		Rectal swab	ESBL- <i>E. coli</i> ^a	P-1112
	December 2015	Ear	<i>Staphylococcus pseudintermedius</i> , <i>Acinetobacter</i> sp., yeast	NA
		Rectal swab	ESBL- <i>E. coli</i> ^a	NA
	May 2016	Ear	Non-specific growth	NA
		Rectal swab	Negative	NA
B	December 2013	Ear	<i>S. pseudintermedius</i> , <i>Acinetobacter baumannii</i>	NA
	April 2015	Ear	NDM- <i>E. coli</i> ; MDR- <i>E. coli</i> (MEM-S)	P-1044, (P-1043) ^b , P-1039
	June 2015	Ear	No growth	NA
	September 2015	Ear	<i>Mucor circinelloides</i> (fungus)	NA
		Rectal swab	ESBL- <i>E. coli</i>	P-1113
	December 2015	Ear	<i>S. pseudintermedius</i> , <i>Corynebacterium</i> sp.	NA
		Rectal swab	Negative	NA
	May 2016	Ear	Non-specific growth	NA
		Rectal swab	Negative	NA

CoNS: coagulase-negative staphylococci; MDR: Multidrug resistant; MEM-S: meropenem susceptible; NA: not applicable (no isolate stored); NDM: New-Delhi metallo-beta-lactamase producing,

^a Identical resistance profiles.

^b P-1043 was identical to P-1044 (antibiogram, pulsed-field gel electrophoresis, multilocus sequence typing), except it had a rough colony type.

72 °C (15 s), and final elongation at 72 °C (60 s). The PCR parameters for *icd*, *mdh*, *purA* and *recA* multiplication were the same, except for an annealing temperature of 59 °C (10 s).

Multiplex PCRs in a Phire Green Hot Start II PCR Master Mix (MM, total volume 20 µL) (Thermo Scientific, US) were performed to screen ESBL (CTX-M, TEM and SHV) and AmpC (CIT, DHA, ACC and FOX) gene families. All primers used in these and in the sequencing of CTX-M, as well as PCR conditions, are presented in Table 1 [19–22]. The reaction mixture for ESBL multiplex PCR and CTX-M sequencing contained 15.4 µL of MM, 0.6 µL of dimethyl sulfoxide (DMSO), 0.25 µM of each primer (Oligomer Oy, Finland) and 1 µL of template DNA. The AmpC multiplex PCR contained 13.8 µL of MM, 0.6 µL of DMSO and 0.25 µM of each primer, except for FOX primers, for which the concentration was 0.4 µM. Carbapenemase gene families (KPC, IMP, VIM, NDM, OXA, GES, and IMI) were detected with multiplex PCR (Table 1) [15]. The reaction mixture contained 0.08 U/µL of AmpliTaq Gold polymerase (Applied Biosystems, US) in a 1x reaction buffer (AmpliTaq Gold buffer 10x Mg free, Applied Biosystems, US), 2.0 mM of MgCl₂, 0.2 mM of nt and 0.3 pmol/µL of each primer. A mixture of DNA from bacterial strains (characterised by THL) carrying either the ESBL (multiplex ESBL PCR), AmpC

(multiplex AmpC PCR) or carbapenemase (multiplex carbapenemase) genes mentioned above were used as positive controls in the respective reactions. PCR-grade water was used as a non-template control in each assay.

For sequencing of MLST genes and CTX-M, the PCR products were purified with ExoI and FastAP (Thermo Scientific, US). Sequencing was performed at Macrogen Inc. (the Netherlands) using an ABI 3730 XL automated sequencer. MLST sequences were analysed using CLC Main Workbench (version 6.9.1, Qiagen, Denmark) and the CLC MLST module (version 1.4.7, Qiagen, Denmark). CTX-M sequencing data were analysed using CLC Main Workbench (version 7.5.1) and compared with the National Center for Biotechnology Information (NCBI) library.

Whole genome sequencing

DNA was extracted using the MagAttract HMW DNA Kit (Qiagen, Germany). Whole genome sequencing (WGS) was performed for carbapenem-resistant isolates using an Illumina MiSeq sequencer (Illumina Inc., US) with the Illumina Nextera XT DNA Sample Preparation Kit, the Nextera XT Index Kit with 24 indices for library preparation and MiSeq Reagent Kit V2 (300 cycles) using 150-bp paired-end sequencing. The reads were compiled

FIGURE 1

Pulsed-field gel electrophoresis cluster analysis of the ESBL- and NDM- *Escherichia coli* strains isolated from two dogs and their owners, Finland, March 2015–September 2016 (n = 10 isolates)

XbaI-PFGE pattern	Origin	Strain ID	Date (mo/yr)	Antibiogram ^a	MLST type	Resistance genotype	BioSample accession
	Dog A	P-1112	09/2015	SSRRSRRSSS	ST69	CTX-M Group 9	NA
	Dog B	P-1113	09/2015	SSRRSRRSSS	ST69	CTX-M Group 9	NA
	Human A	O-58	05/2015	SSRRSRRSSS	ST69	CTX-M Group 9	NA
	Human B	O-60	05/2015	SSRRSRRSSS	ST69	CTX-M Group 9	NA
	Dog A	P-1009	03/2015	RRRRSSRRRR	ST167	NDM-5, CMY-42, TEM-1B	SAMN08329266
	Dog A	P-1036	04/2015	RRRRSSRRRR	ST167	NDM-5, CMY-42, TEM-1B	SAMN08329265
	Dog B	P-1039 ^b	04/2015	SSRRSRRSSR	ST167	CMY-42	SAMN08329267
	Dog B	P-1043 ^c	04/2015	RRRRSSRRRR	ST167	NDM-5, CMY-42, TEM-1B	SAMN08329268
	Dog B	P-1044 ^c	04/2015	RRRRSSRRRR	ST167	NDM-5, CMY-42, TEM-1B	SAMN08329269
	Human A	O-59	05/2015	RRRRSSRRRR	ST167	NDM-5, CMY-42	SAMN08329270

ESBL: extended-spectrum beta-lactamase; MLST: multilocus sequence typing; NA: not available; NDM: New-Delhi metallo-beta-lactamase producing.

^a Antibiogram is represented in the following order: amikacin, amoxicillin-clavulanic acid, ampicillin, cefpodoxime, chloramphenicol, doxycycline, enrofloxacin, gentamicin, meropenem, and trimethoprim-sulfamethoxazole.

^b P-1039 is meropenem susceptible MDR-*E. coli*.

^c These two strains were otherwise identical, but P-1043 had a rough colony type.

using Velvet assembler version 1.1.04 included in Ridom SeqSphere+ software (Ridom SeqSphere+ version 2.4.0; Ridom GmbH, Germany). Beta-lactamase resistance genes were detected using ResFinder 2.1 on the Center for Genomic Epidemiology (CGE) server [23]. Plasmids were detected using PlasmidFinder 1.3 [24].

An *ad hoc* core-genome MLST (cgMLST) protocol for *E. coli*, containing 2,634 target alleles, was used on SeqSphere+ to compare the strains originating from the family (i.e. the dogs and owners) to strains previously isolated from human patients in Finland, with similar ST and carbapenemase profile, which were identified from the National Infectious Diseases Register at THL. The *ad hoc* cgMLST scheme was created using the cgMLST Target Definer of Ridom SeqSphere software and genome sequences obtained from GenBank. The whole-genome sequence of *Escherichia coli* strain G150 (GenBank accession number: LQHK01000008) was used as a reference strain and three *E. coli* strains, TW14588 (GenBank accession number: NZ_CM000662), 453 (GenBank accession number: NZ_MPGRo1000001) and 8368 (GenBank accession number: NZ_CP017444), were used as query genomes.

Results

Case presentations

The patients were two male Finnish Hound dogs, Dog A (five years-old) and Dog B (two-years old), living in the same family. Both dogs had a long history of recurrent *otitis externa*, the underlying factor of which was unidentified, although allergy/atopy was suspected. Dog A had its first *otitis externa* episode in

February 2013, when a local veterinarian diagnosed ear inflammation with yeast overgrowth in the dog's right ear. Dog B began experiencing ear problems in October 2013. Both dogs continued to experience intermittent symptoms of ear infection, despite both topical and systemic antimicrobial (Table 2) and non-antimicrobial therapy, including corticosteroids.

Otitis media was diagnosed in the left ear of Dog A in November 2014, while the same condition had been diagnosed in the left ear of Dog B in December 2013. In March 2015, a bacteriological specimen from the left ear of Dog A was sent to the CML. The specimen revealed a meropenem-resistant MDR *E. coli* (for resistance profile, see Figure 1), but carbapenemase production, according to the modified Hodge's test, appeared to be negative. At that time, the *E. coli* isolate was preserved for further investigation due to the unusual resistance profile. The veterinarian was informed about the unusual phenotype and was advised to handle the dog with contact isolation precautions. Specimens from the ears of both dogs were taken in late April 2015. The specimen from Dog A again revealed meropenem-resistant MDR *E. coli*, along with non-specific growth. This *E. coli* isolate gave a positive result for carbapenemase production in the modified Hodge's test. This isolate and the earlier meropenem-resistant *E. coli* from Dog A were both metallo-beta-lactamase positive in the double-disk diffusion test. Dog B's specimen initially revealed only MDR *E. coli* that was meropenem sensitive, but after the meropenem-resistant *E. coli* finding from Dog A, Dog B's specimen was re-cultured onto a chromogenic agar (UriSelect, Bio-Rad, US) on which a meropenem disc was placed. This revealed a

FIGURE 2

Relatedness of NDM-5 *Escherichia coli* ST167 isolates based on cgMLST analysis, Finland, 2014–2015

cgMLST: core genome multilocus sequence typing; NDM: New-Delhi metallo-beta-lactamase producing.

Minimum spanning tree based on comparing of 2,634 target genes and ignoring missing values in pairwise comparison. The dashed circle defines the strains of this study. Strains H1 to H4 are NDM-5 positive *E. coli* ST167 strains from humans isolated in 2014 and 2015. The numbers above the lines connecting the strains indicate allelic differences. P-1036 and P-1009 are from dog A; P-1044, P-1043, and P-1039 are from Dog B (P-1043 was isolated together with P-1044, was otherwise similar but had a rough colony type). O-59 is from human A. All these were NDM-5 positive except for P-1039.

meropenem-resistant *E. coli* isolate with the same anti-biogram as Dog A's isolates. Systemic treatment with antimicrobials was stopped and reasonable control of the ear infection in both dogs was achieved by topical chloramphenicol treatment and anti-inflammatory therapy. The specimen histories of the dogs with respective findings are presented in Table 3.

Epidemiological investigation

The dogs were domestic Finnish Hounds from the same family in rural Eastern Finland. The family consisted of only two adults, and the only animals were the dogs in question. Both family members were of Finnish ethnicity. The dogs lived together in a fenced yard with a dog house and were used for hunting hares. Their diet mainly comprised hare and elk offal, in addition to raw commercial dog food (domestic). They also received cooked cow spines (domestic) and, occasionally, leather bones (unknown origin). The dogs were taken to the woods for hunting or training approximately three times per week. The dogs had not travelled outside of Finland and had not been in contact with other animals while at home. However, while hunting, they had contact with wild animals, mainly hares, but probably also with mice and voles. The human family members had visited Estonia, Sweden, Norway and Croatia within the previous 2 years. Their medical history included hospitalisations in Finland, but not abroad. Neither of the dog owners worked in the healthcare sector.

Bacteriological and molecular investigations

The two human family members were screened on a voluntary basis for the presence of carbapenemase/ESBL-producing Gram-negative rods. One of them carried NDM- as well as ESBL-producing *E. coli*, while the other family member only carried ESBL-producing *E. coli*. Altogether, seven canine and three human *E. coli* isolates were further typed. One isolate (P-1043) from Dog B was identical to isolate P-1044 from the same dog, apart from colony morphology (Figure 1). All meropenem-resistant isolates ($n=5$) belonged to ST167, had an identical antibiogram and PFGE profile, showed metallo-beta-lactamase activity, and were positive for NDM, CIT and TEM by multiplex PCR, except for the human isolate O-59, which was TEM-negative. In addition, one canine isolate susceptible to meropenem and amikacin belonged to the same cluster. All ESBL *E. coli* isolates ($n=4$) were part of the same PFGE cluster and MLST type (ST69), with a characteristic antibiogram. Sequencing confirmed the presence of a CTX-M group 9 gene in the isolates of the latter cluster (Figure 1).

WGS was performed on five canine ST167 *E. coli* isolates and the human NDM-positive ST167 isolate. Sequences were submitted to the NCBI BioSample database (see Figure 1 for accession numbers). All five meropenem-resistant isolates carried NDM-5. All six isolates had CMY-42 in their WGS profile (Figure 1).

Ad hoc cgMLST grouped the six isolates very tightly: one strain from Dog A and the owner's strain were identical, as were two strains from Dog B. Other strains, including one without NDM-5, had only 1 to 2 allelic differences (Figure 2). The search for previous isolates in Finland of the same type and antibiogram as the ones found in the family revealed four human *E. coli* ST167 isolates, all NDM-5 positive, from 2014–15. In the cgMLST analysis these isolates were clearly different from the strains of this study, differing in at least 67 alleles. All the NDM-5-positive *E. coli* strains of this study had *IncI1*- and *IncFII*-type plasmids, while the strain without the NDM-5 gene had only *IncI1*-type plasmids.

A retrospective search of the CML database yielded CTX-M-positive *E. coli* ST167 isolates in horses in Finland, but these were vastly different in PFGE compared with the NDM-5 *E. coli* of this study (data not shown). No other previous *E. coli* ST69 of animal origin was observed.

Follow-up specimens

Carbapenemase-producing *E. coli* was not detected in canine follow-up specimens, but rectal screenings of both dogs in September 2015 yielded CTX-M-positive *E. coli* ST69 (Table 3). Only one follow-up specimen was taken from human A, who originally carried both NDM and CTX-M-positive *E. coli* strains. The follow-up specimen again yielded ESBL *E. coli* with a similar antibiogram to the other ST69 isolates in this study. However, this isolate was not available for typing.

Discussion

The results from molecular analyses strongly indicate the transmission of ST167 NDM-5 and ST69 CTX-M group 9 *E. coli* between two dogs and humans in the same family. The transfer of ESBL-producing *Enterobacteriaceae* between humans and dogs has previously been described [25], but to our best knowledge, this is the first report on the transmission of carbapenemase-producing *E. coli* between dogs and humans.

We consider that the transmission of ST167 NDM-5 had most likely occurred from human to dog. There are several reasons for this. Firstly, CPE are more frequent in humans than in animals [6]. Secondly, according to resistance surveillance, carbapenem-resistant bacteria have never been observed in animals in Finland before this report. They have, however, been observed with a gradually increasing frequency in humans, most of whom probably acquired the bacteria abroad [26]. In 2008–15, among 31 Finnish human NDM cases, 11 were NDM-5 *E. coli*. Seven of these had been imported from India, and one each from Croatia, Thailand and Egypt, while one was of unknown origin [26]. The human family members of this study had a history of hospitalisation in Finland, but not abroad. They also had visited Croatia in May 2013, which might be one possible origin for the NDM-5 *E. coli* ST167. The Balkan region has

been reported as a potential reservoir of NDM genes, particularly NDM-1 [27]. It is also unlikely that the dogs had acquired NDM *E. coli* from their feed, although it is known that exposure to raw meat increases the risk of ESBL carriage in dogs [28,29]. However, carbapenem-resistant isolates have not been detected in food-producing animals in Finland [30], and even the presence of ESBL/AmpC-type beta-lactamases (CTX-M-1 and CMY-2) is deemed to be at a low level in Finnish livestock (0.8%, 1% and 8.1% in cattle, pigs and broilers, respectively in 2013–14) [31]. No data exist on the prevalence of CPE among wild animals or in the environment in Finland, and these cannot therefore be ruled out as potential sources of infection for the dogs or the owners. However, the general level of resistance in Finnish wild animals was studied in 1999, and was found to be much lower than in the human population [32].

The origin of the CTX-M-9 *E. coli* ST69 isolate is unclear. So far, ST69 *E. coli* have not been observed in companion animals in Finland (data not shown), while CTX-M is a common ESBL family among canine ESBL isolates [33]. However, no data are yet available on the genetic variants of canine CTX-M in Finland.

Reports concerning carbapenem-resistant bacteria in companion animals are uncommon [34–42]. In 2013, a US study reported six unrelated NDM *E. coli* isolates collected in 2008–2009 from five canine and one feline patient, of which four were confirmed to be NDM-1 [39]. This is interesting, since the first report of NDM-1 carbapenemase in a *K. pneumoniae* and an *E. coli* isolate from a Swedish human patient of Indian origin was observed in 2008 and published a year later [43]. The first NDM-5-positive *E. coli* of animal origin was reported in an Algerian dog in 2015 [42]. The sequence type of the NDM-5 *E. coli* was ST1284, which is a double locus variant of *E. coli* ST167. In addition, OXA-48 has been observed in *E. coli* in companion animals (dogs and cats) in Algeria (2014–15) [37], the US (2009–13) [35] and France (dogs) (2015) [40], and in *E. coli* and *K. pneumoniae* at a veterinary clinic in Germany (dogs) (2012) [41]. Furthermore, OXA-23 *Acinetobacter* has been reported in dogs in France (2012) [36], VIM-1-producing *K. pneumoniae* in a dog in Spain (2014–15) [38], and IMP-4 *Salmonella* Typhimurium in cats at an animal shelter in Australia (year not specified) [34]. Carbapenem-producing bacteria have also been reported in horses and food-producing animals (NDM in poultry and pigs, OXA-23 in dairy cows), as well as in environmental specimens (2003–13) [6].

It seems that the presence of carbapenemase-producing bacteria in animals is not linked to carbapenem use, since carbapenem products are not authorised for veterinary use [6]. Some off-label use could exist in companion animals according to the cascade principle (Articles 10 and 11 of Directive 2001/82/EC) in the European Union [44], but the extent of such use is unknown. In Finland, however, national legislation prohibits the use of carbapenems in veterinary

medicine (Government decree 1054/2014). Our investigation did not reveal any carbapenem use in the dogs of this study, but the selection pressure caused by numerous other antimicrobials was probably enough to co-select carbapenem resistance after the acquisition of carbapenem-resistant MDR *E. coli*. For example, ESBL-producing *Enterobacteriaceae* may be co-selected for by fluoroquinolone therapy [45]. Both dogs had received local as well as systemic fluoroquinolone therapy, which may have contributed to the survival of the MDR NDM-5 *E. coli*.

The NDM-5 *E. coli* isolates in our study belonged to ST167. A database search revealed that only four ST167 NDM-5-positive *E. coli* strains had previously been isolated from humans in Finland. In three cases, the patients had had contact abroad, i.e. Thailand, India and Egypt. The results of cgMLST analyses revealed that previous NDM-5 ST167 *E. coli* were not related to the cluster of this study. Previously, the dissemination of NDM-5 has been linked to *IncX3* plasmids [46], while the *IncI1* and *IncFII* plasmids that were present in the NDM-5 isolates of this study have been associated with NDM-1 isolates [47]. *IncFII* was also identified from all previously isolated NDM-5 ST167 *E. coli* strains that were investigated, although these were not related to the strains from the cases presented here. It is possible that all NDM-5-positive dog-cluster *E. coli* strains carried the NDM-5 gene in *IncFII*-type plasmids, and the one carbapenem-susceptible strain without NDM-5 had lost this plasmid.

As carbapenemase-producing strains and carbapenem resistance genes spread around the world due to travel and the international trade of food and feed, it is likely that such bacteria will be detected in animals in increasing numbers [5], as has been the case with ESBL/AmpC *Enterobacteriaceae*. Close contact with companion animals increases the likelihood of human-to-animal CPE transmission. Once it has entered the animal population, a CPE can be efficiently co-selected by other antimicrobial classes due to the MDR character of CPE. The widespread use of antimicrobials in companion animals could thus enable them to act as reservoirs for CPE isolates and other resistant bacteria. In our cases, all systemic antimicrobials were discontinued and only topical therapy was used. This may have facilitated the reduction of NDM-5 and ESBL *E. coli* below the level of detection in follow-up specimens.

An enrichment method for screening would have been more sensitive, but was not used, as it is not recommended by the national guidelines. Furthermore, the use of the European Committee on Antimicrobial Susceptibility Testing (EUCAST) epidemiological cut-off value (ECOFF) for meropenem (<25 mm) [48] instead of the CLSI meropenem screening breakpoint (≤22 mm) might have increased the likelihood of detecting non-wild-type isolates. However, as a clinical laboratory, adjusting breakpoints for this specific case was

not considered necessary. Moreover, the meropenem histogram from *E. coli* isolates of the CML revealed that the CLSI breakpoint is capable of satisfactorily differentiating non-wild-type isolates from the wild-type population.

In conclusion, to our knowledge, these are the first canine clinical NDM-5 *E. coli* findings in Europe and this is the first confirmed case of transmission between dogs and humans. While the original source remains unclear, it is likely that either one or both of the dogs had acquired the strain from a human source. Although carbapenems had not been administered to these dogs, frequent exposure to diverse systemic antimicrobials probably facilitated the propagation of the bacteria in the canine host. It is vital for veterinary laboratories to remain alert, and to screen *Enterobacteriaceae* isolates for carbapenem and third-generation cephalosporin resistance. This could be done by using suitable screening disks in testing panels or by using, for example, commercial screening plates for preliminary screening. Isolates with reduced susceptibility to carbapenems should be sent to a reference laboratory for confirmation. Furthermore, the veterinary community should continue to emphasise the importance of prudent antimicrobial use.

Acknowledgements

The authors wish to thank the owner of the dogs and the treating veterinarian for their excellent cooperation. We also thank laboratory personnel at the University of Helsinki, the Institute of Health and Welfare, and the Food Safety Authority Evira for their invaluable assistance. The Alfred Kordelin Foundation, the Finnish Veterinary Foundation, the Foundation for Canine Health Research, and the Finnish Foundation of Veterinary Research, the Finnish Cultural Foundation are thanked for their financial support. Roy Siddall is acknowledged for revising the language of the revised manuscript.

Conflict of interest

None declared.

Authors' contributions

MR and TG performed identification and susceptibility testing, including ESBL phenotype and modified Hodge testing of the bacterial isolates. MR performed the CML database search for previous isolates. MR supervised and TG performed the epidemiological investigation, MÖ and JJ performed carbapenemase gene PCR, imipenem hydrolysis testing, WGS analysis and THL database search for previous isolates, ME performed PFGE, MLST and PCR for ESBL, SN and KP performed double disk diffusion for metallo-beta-lactamase testing. All authors participated in the drafting the manuscript, and reviewed and approved the final version of the manuscript.

References

1. Shaikh S, Fatima J, Shakil S, Rizvi SM, Kamal MA. Antibiotic resistance and extended spectrum beta-lactamases: Types,

- epidemiology and treatment. *Saudi J Biol Sci.* 2015;22(1):90-101. <https://doi.org/10.1016/j.sjbs.2014.08.002> PMID: 25561890
2. Meletis G. Carbapenem resistance: overview of the problem and future perspectives. *Ther Adv Infect Dis.* 2016;3(1):15-21. <https://doi.org/10.1177/2049936115621709> PMID: 26862399
3. Wang Q, Zhang Y, Yao X, Xian H, Liu Y, Li H, et al. Risk factors and clinical outcomes for carbapenem-resistant Enterobacteriaceae nosocomial infections. *Eur J Clin Microbiol Infect Dis.* 2016;35(10):1679-89. <https://doi.org/10.1007/s10096-016-2710-0> PMID: 27401905
4. Lee CR, Lee JH, Park KS, Kim YB, Jeong BC, Lee SH. Global Dissemination of Carbapenemase-Producing *Klebsiella pneumoniae*: Epidemiology, Genetic Context, Treatment Options, and Detection Methods. *Front Microbiol.* 2016;7:895. <https://doi.org/10.3389/fmicb.2016.00895> PMID: 27379038
5. Guerra B, Fischer J, Helmuth R. An emerging public health problem: acquired carbapenemase-producing microorganisms are present in food-producing animals, their environment, companion animals and wild birds. *Vet Microbiol.* 2014;171(3-4):290-7. <https://doi.org/10.1016/j.vetmic.2014.02.001> PMID: 24629777
6. Woodford N, Wareham DW, Guerra B, Teale C. Carbapenemase-producing Enterobacteriaceae and non-Enterobacteriaceae from animals and the environment: an emerging public health risk of our own making? *J Antimicrob Chemother.* 2014;69(2):287-91. <https://doi.org/10.1093/jac/dkt392> PMID: 24092657
7. McLaughlin M, Advincula MR, Malczynski M, Qi C, Bolon M, Scheetz MH. Correlations of antibiotic use and carbapenem resistance in enterobacteriaceae. *Antimicrob Agents Chemother.* 2013;57(10):5131-3. <https://doi.org/10.1128/AAC.00607-13> PMID: 23836188
8. Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect.* 2012;18(3):268-81. <https://doi.org/10.1111/j.1469-0691.2011.03570.x> PMID: 21793988
9. Valtioneuvoston asetus eräiden lääkeaineiden käytön kieltämisestä eläimille. [Government Decree concerning the prohibition of the use of certain veterinary medicinal products for animals]. Finlex Data Bank. Finnish. Available from: <https://www.finlex.fi/fi/laki/alkup/2014/20141054>
10. National Institute for Health and Welfare (THL). Tartuntataudit Suomessa 2016. [Infectious diseases in Finland 2016]. Helsinki, Finland: THL; 2017. Report No.: 10/2015. Finnish.
11. Garcia LS, Isenberg HD. Clinical microbiology procedures handbook. 3rd ed. Washington, DC: ASM Press; 2010.
12. Clinical and Laboratory Standards Institute (CLSI). Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated From Animals: Approved Standard VET01-A4. Wayne, PA: CLSI; 2013.
13. Clinical and Laboratory Standards Institute (CLSI). Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated From Animals: Second Informational Supplement VET01-S2. Wayne, PA: CLSI; 2013.
14. Clinical and Laboratory Standards Institute (CLSI). Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Fourth Informational Supplement M100-S24. Wayne, PA: CLSI; 2014.
15. Österblad M, Kirveskari J, Hakanen AJ, Tissari P, Vaara M, Jalava J. Carbapenemase-producing Enterobacteriaceae in Finland: the first years (2008-11). *J Antimicrob Chemother.* 2012;67(12):2860-4. <https://doi.org/10.1093/jac/dks299> PMID: 22855858
16. National Institute for Health and Welfare (THL). Ohje moniresistenttien mikrobien tartunnantorjunnasta. [Manual on infection prevention for multi-drug resistant microbes]. Helsinki, Finland: THL; 2017. Report No.: 22/2017. Finnish.
17. Center for Disease Control and Prevention (CDC). One-Day (24-28 h) Standardized Laboratory Protocol for Molecular Subtyping of *Escherichia coli* O157:H7, non-typhoidal *Salmonella* serotypes, and *Shigella sonnei* by Pulsed Field Gel Electrophoresis (PFGE). Wayne, PA: CDC, PulseNet; 2013.
18. Lau SH, Reddy S, Cheesbrough J, Bolton FJ, Willshaw G, Cheasty T, et al. Major uropathogenic *Escherichia coli* strain isolated in the northwest of England identified by multilocus sequence typing. *J Clin Microbiol.* 2008;46(3):1076-80. <https://doi.org/10.1128/JCM.02065-07> PMID: 18199778
19. Mulvey MR, Soule G, Boyd D, Demczuk W, Ahmed RM. Multi-provincial *Salmonella* Typhimurium Case Control Study Group. Characterization of the first extended-spectrum beta-lactamase-producing *Salmonella* isolate identified in Canada. *J Clin Microbiol.* 2003;41(1):460-2. <https://doi.org/10.1128/JCM.41.1.460-462.2003> PMID: 12517894
20. Nyberg SD, Osterblad M, Hakanen AJ, Huovinen P, Jalava J. The Finnish Study Group For Antimicrobial Resistance. Detection and molecular genetics of extended-spectrum beta-lactamases among cefuroxime-resistant *Escherichia coli* and *Klebsiella* spp. isolates from Finland, 2002-2004. *Scand J Infect Dis.* 2007;39(5):417-24. <https://doi.org/10.1080/00365540601105731> PMID: 17464864
21. Haanperä M, Forssten SD, Huovinen P, Jalava J. Typing of SHV extended-spectrum beta-lactamases by pyrosequencing in *Klebsiella pneumoniae* strains with chromosomal SHV beta-lactamase. *Antimicrob Agents Chemother.* 2008;52(7):2632-5. <https://doi.org/10.1128/AAC.01259-07> PMID: 18458132
22. Pérez-Pérez FJ, Hanson ND. Detection of plasmid-mediated AmpC beta-lactamase genes in clinical isolates by using multiplex PCR. *J Clin Microbiol.* 2002;40(6):2153-62. <https://doi.org/10.1128/JCM.40.6.2153-2162.2002> PMID: 12037080
23. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother.* 2012;67(11):2640-4. <https://doi.org/10.1093/jac/dks261> PMID: 22782487
24. Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, et al. In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother.* 2014;58(7):3895-903. <https://doi.org/10.1128/AAC.02412-14> PMID: 24777092
25. Ljungquist O, Ljungquist D, Myrenäs M, Rydén C, Finn M, Bengtsson B. Evidence of household transfer of ESBL-/pAmpC-producing Enterobacteriaceae between humans and dogs - a pilot study. *Infect Ecol Epidemiol.* 2016;6(1):31514. <https://doi.org/10.3402/iee.v6.31514> PMID: 27330043
26. National Institute for Health and Welfare (THL). Tartuntataudit Suomessa 2015. [Infectious diseases in Finland 2015]. Helsinki, Finland: THL; 2016. Report No.: 10/2015. Finnish.
27. Berrazeg M, Diene S, Medjahed L, Parola P, Drissi M, Raoult D, et al. New Delhi Metallo-beta-lactamase around the world: an eReview using Google Maps. *Euro Surveill.* 2014;19(20):20809. <https://doi.org/10.2807/1560-7917.ES2014.19.20.20809> PMID: 24871756
28. Nilsson O. Hygiene quality and presence of ESBL-producing *Escherichia coli* in raw food diets for dogs. *Infect Ecol Epidemiol.* 2015;5(1):28758. <https://doi.org/10.3402/iee.v5.28758> PMID: 26490763
29. Baede VO, Wagenaar JA, Broens EM, Duim B, Dohmen W, Nijssen R, et al. Longitudinal study of extended-spectrum-beta-lactamase- and AmpC-producing Enterobacteriaceae in household dogs. *Antimicrob Agents Chemother.* 2015;59(6):3117-24. <https://doi.org/10.1128/AAC.04576-14> PMID: 25779568
30. FINRES-Vet. FINRES-Vet 2013-2015 Finnish Veterinary Antimicrobial Resistance Monitoring and Consumption of Antimicrobial Agents. Finnish Food Safety Authority Evira; 2017. 68 p. Available from: www.evira.fi
31. Päivärinta M, Pohjola L, Fredriksson-Ahomaa M, Heikinheimo A. Low Occurrence of Extended-Spectrum beta-lactamase-Producing *Escherichia coli* in Finnish Food-Producing Animals. *Zoonoses Public Health.* 2016;63(8):624-31. <https://doi.org/10.1111/zph.12277> PMID: 27215423
32. Österblad M, Norrdahl K, Korpimäki E, Huovinen P. Antibiotic resistance. How wild are wild mammals? *Nature.* 2001;409(6816):37-8. <https://doi.org/10.1038/35051173> PMID: 11343104
33. Jalava JVN, Miettinen S, Pelkonen S, Rantala M. Prevalence of third-generation cephalosporin-resistant *Escherichia coli* and their resistance mechanisms in dogs in Finland, P1107 abstract. 22nd ECCMID Conference; 31 March – 3 April 2012; London, UK; 2012.
34. Abraham S, O'Dea M, Trott DJ, Abraham RJ, Hughes D, Pang S, et al. Isolation and plasmid characterization of carbapenemase (IMP-4) producing *Salmonella enterica* Typhimurium from cats. *Sci Rep.* 2016;6(1):35527. <https://doi.org/10.1038/srep35527> PMID: 27767038
35. Liu X, Thungrat K, Boothe DM. Occurrence of OXA-48 Carbapenemase and Other beta-Lactamase Genes in ESBL-Producing Multidrug Resistant *Escherichia coli* from Dogs and Cats in the United States, 2009-2013. *Front Microbiol.* 2016;7:1057. PMID: 27462301
36. Hérivaux A, Pailhoriès H, Quinqueneau C, Lemarié C, Joly-Guillou ML, Ruvoen N, et al. First report of carbapenemase-producing *Acinetobacter baumannii* carriage in pets from the community in France. *Int J Antimicrob Agents.* 2016;48(2):220-1. <https://doi.org/10.1016/j.ijantimicag.2016.03.012> PMID: 27424597
37. Yousfi M, Touati A, Mairi A, Brasme L, Gharout-Sait A, Guillard T, et al. Emergence of Carbapenemase-Producing *Escherichia coli* Isolated from Companion Animals in Algeria. *Microb Drug Resist.* 2016;22(4):342-6. <https://doi.org/10.1089/mdr.2015.0196> PMID: 26741510

38. González-Torralba A, Oteo J, Asenjo A, Bautista V, Fuentes E, Alós JL. Survey of Carbapenemase-Producing Enterobacteriaceae in Companion Dogs in Madrid, Spain. *Antimicrob Agents Chemother.* 2016;60(4):2499-501. <https://doi.org/10.1128/AAC.02383-15> PMID: 26824947
39. Shaheen BW, Nayak R, Boothe DM. Emergence of a New Delhi metallo- β -lactamase (NDM-1)-encoding gene in clinical *Escherichia coli* isolates recovered from companion animals in the United States. *Antimicrob Agents Chemother.* 2013;57(6):2902-3. <https://doi.org/10.1128/AAC.02028-12> PMID: 23587948
40. Melo LC, Boisson MN, Saras E, Médaille C, Boulouis HJ, Madec JY, et al. OXA-48-producing ST372 *Escherichia coli* in a French dog. *J Antimicrob Chemother.* 2017;72(4):1256-8. PMID: 28039279
41. Stolle I, Prenger-Berninghoff E, Stamm I, Scheufen S, Hassdenteufel E, Guenther S, et al. Emergence of OXA-48 carbapenemase-producing *Escherichia coli* and *Klebsiella pneumoniae* in dogs. *J Antimicrob Chemother.* 2013;68(12):2802-8. <https://doi.org/10.1093/jac/dkt259> PMID: 23833179
42. Yousfi M, Mairi A, Bakour S, Touati A, Hassissen L, Hadjadj L, et al. First report of NDM-5-producing *Escherichia coli* ST1284 isolated from dog in Bejaia, Algeria. *New Microbes New Infect.* 2015;8:17-8. <https://doi.org/10.1016/j.nmni.2015.09.002> PMID: 26566444
43. Yong D, Toleman MA, Giske CG, Cho HS, Sundman K, Lee K, et al. Characterization of a new metallo-beta-lactamase gene, bla(NDM-1), and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India. *Antimicrob Agents Chemother.* 2009;53(12):5046-54. <https://doi.org/10.1128/AAC.00774-09> PMID: 19770275
44. Directive 2001/82/EC of the European Parliament and of the Council of 6 November 2001 on the Community code relating to veterinary medicinal products. *OJ L* 311, 28.11.2001, p. 1–66. Available from: <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=celex%3A32001L0082>
45. Kantele A, Mero S, Kirveskari J, Lääveri T. Fluoroquinolone antibiotic users select fluoroquinolone-resistant ESBL-producing Enterobacteriaceae (ESBL-PE) - Data of a prospective traveller study. *Travel Med Infect Dis.* 2017;16:23-30. <https://doi.org/10.1016/j.tmaid.2017.01.003> PMID: 28153711
46. Yang P, Xie Y, Feng P, Zong Z. blaNDM-5 carried by an IncX3 plasmid in *Escherichia coli* sequence type 167. *Antimicrob Agents Chemother.* 2014;58(12):7548-52. <https://doi.org/10.1128/AAC.03911-14> PMID: 25246393
47. Kumarasamy KK, Toleman MA, Walsh TR, Bagaria J, Butt F, Balakrishnan R, et al. Emergence of a new antibiotic resistance mechanism in India, Pakistan, and the UK: a molecular, biological, and epidemiological study. *Lancet Infect Dis.* 2010;10(9):597-602. [https://doi.org/10.1016/S1473-3099\(10\)70143-2](https://doi.org/10.1016/S1473-3099(10)70143-2) PMID: 20705517
48. European Committee on Antimicrobial Susceptibility Testing (EUCAST). Antimicrobial wild type distributions of microorganisms: The European Committee on Antimicrobial Susceptibility Testing; Växjö: EUCAST; 2017. Available from: <https://mic.euCAST.org/Eucast2/>

License and copyright

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0) Licence. You may share and adapt the material, but must give appropriate credit to the source, provide a link to the licence, and indicate if changes were made.

This article is copyright of the authors, 2018.