Detection of *Aedes flavopictus* (Yamada, 1921), Netherlands, June 2019

Adolfo Ibáñez-Justicia¹, Bart van de Vossenber², Rens van den Biggelaar¹, Joris Voogd², Eveline Metz², Frans Jacobs¹, Marian Dik¹, Arjan Stroo¹

1. Centre for Monitoring of Vectors, National Reference Centre, Netherlands Food and Consumer Product Safety Authority, Wageningen, Netherlands

2. Molecular Biology Group, National Reference Centre, Netherlands Food and Consumer Product Safety Authority, Wageningen, Netherlands

Correspondence: Adolfo Ibáñez-Justicia (a.ibanezjusticia@nvwa.nl)

Citation style for this article:

Ibáñez-Justicia Adolfo, van de Vossenber Bart, van den Biggelaar Rens, Voogd Joris, Metz Eveline, Jacobs Frans, Dik Marian, Stroo Arjan. Detection of *Aedes flavopictus* (Yamada, 1921), Netherlands, June 2019. Euro Surveill. 2019;24(30):pii=1900433. <https://doi.org/10.2807/1560-7917.ES.2019.24.30.1900433>

Article submitted on 04 Jul 2019 / accepted on 25 Jul 2019 / published on 25 Jul 2019

In June 2019, a single specimen collected at a used tyre company was identified as *Aedes flavopictus* (Yamada, 1921), a sibling species of *Ae. albopictus*. *Ae. flavopictus* has not been recorded outside Japan and South Korea. Although it has only shown dengue virus vector competence under laboratory conditions, its detection demonstrates the value of active surveillance at risk locations and molecular tools for timely intervention against exotic mosquitoes with potential future public health impact.

On 13 June 2019, in a sample of mosquitoes captured in the province of Flevoland, the Netherlands during routine surveillance, a specimen was recognised as potentially belonging to an exotic species. Based on the molecular analysis of the specimen, we concluded that it belonged to *Aedes flavopictus* (Yamada, 1921). To our knowledge, this is both the first time that this species has been reported outside its area of origin in north-east Asia and associated with the import of used tyres.

Mosquito surveillance at high-risk locations

From 2010 onwards, an annual quantitative risk assessment of all used tyre companies known to import tyres into the Netherlands is performed by the Centre for Monitoring of Vectors (CMV) of the National Reference Centre (NRC) of the Netherlands Food and Consumer Product Safety Authority (NVWA) for the purpose of assessing the risk of invasive mosquito species (IMS) introduction. Sampling methods and frequency of the inspections depend on the expected risk of that location [1]. A company in the province of Flevoland has been categorised as high risk based on earlier introductions of *Ae. albopictus* and their international trade with companies in areas where IMS are established. Consequently, mosquito trap sampling is carried out every 2 weeks during the period in which mosquitoes in the Netherlands are most active, i.e. from mid-April to

mid-October. At this location, at least one BG-Sentinel trap lured with BG-Lure (Biogents, Regensburg, Germany) and CO₂ is collecting mosquito specimens 24 hours a day, 7 days a week. Samples collected from used tyre companies are labelled in the field with a unique code, sealed and sent to the vector specialists of the CMV for further analysis.

Detection of *Aedes flavopictus*

At the laboratory, mosquitoes are counted and morphologically identified using the electronic key of Schaffner et al. [2] and the key of Becker et al. [3]. Samples containing only indigenous mosquito species are considered negative samples and samples containing IMS are considered positive samples. In one sample of mosquitoes captured during surveillance at the tyre company in Flevoland, one specimen was recognised as a potentially exotic species. This sample was obtained from the BG-Sentinel trap within 28 May 2019 and 12 June 2019. The sample also contained 138 indigenous *Culex* spp. mosquitoes. All mosquitoes collected were females. The potential exotic specimen was damaged and we were not able to reliably morphologically characterise it to the species level. As it was recognised as an *Aedes* species, one leg of the specimen was subjected to molecular identification using four real-time PCRs specifically developed for identification of *Ae. albopictus*, *Ae. aegypti*, *Ae. atropalpus* and *Ae. japonicus* [4]. All four simplex assays produced negative results and the DNA extracted from the specimen was further subjected to sequencing of the partial mitochondrial *cox1* gene (GenBank accession number MN102102), also referred to as DNA barcoding [5]. Blast analysis of the obtained 658 nt sequence in the Barcode of Life Data System version 4 using the species level barcode records for animal identification (BOLD Systems, Guelph, Canada) and the National Center for Biotechnology Information's (NCBI) GenBank showed 98% coverage and 99.4% pairwise

identity with an accession submitted by Maekawa et al. 2016 (GenBank accession number LCo54355.1) [6]. Clustering of sequence MN102102 with the blast hits from BOLD Systems and NCBI showed grouping of the sequence in an *Ae. flavopictus* species-specific clade. Following the finding of this species, the trap was checked at the site during two additional sampling periods, 12–19 June 2019 and 19–26 June 2019, but only indigenous *Culex* spp. were identified.

Climate suitability assessment

For a rapid assessment of the suitability of the Dutch climate for *Ae. flavopictus*, we used the Match Climates function of the CLIMEX software version 3 (Hearne Scientific Software, Melbourne, Australia). This function is a tool for comparing the meteorological data of different places without reference to any particular species. This function allows us to select a location, the Home location, and find other locations, the 'Away' locations, with climates similar to that of the 'Home' location. In CLIMEX, the similarity of the climate is measured by the Composite Match Index (CMI), a value between zero and one, with higher values corresponding to a greater match between locations [7]. For our study, we used two 'Home' locations: (i) Miyako, the nearest climate station to the location of the specimen that showed 99.4% sequence homology with our specimen (latitude/longitude=40.8523 N/140.5310 E, North of Aomori province in Japan), and (ii) Sapporo, which is on the island of Hokkaido, where the species is known to occur (BOLD accession number GBDCU179–12). Results of the match climates function projected to Europe showed that the CMI values of several locations in the Netherlands, Amsterdam, Rotterdam and Groningen were around 0.7. In CLIMEX, a CMI>0.7 means good matching. It is anticipated that winter temperatures in the Netherlands would not likely be a limiting factor for the species since the averaged minimum temperatures in Sapporo, Hokkaido, Japan are lower than in the Netherlands. This preliminary assessment shows that the climatological conditions are unlikely to be a limiting factor for *Ae. flavopictus* in the Netherlands or other temperate areas in Europe. Other ecological restraints that limit the invasiveness of this species might apply, so further studies are necessary.

Control measures

The origin of the found specimen is currently unknown, and for this reason, the NVWA is performing an intensive survey to find its source.

Notwithstanding the relative low demonstrated risk for public health, the Dutch authorities have put a plan to prevent possible spread and establishment of the species into place. First, the complete company site is being treated by spraying *Bacillus thuringiensis israelensis* (Bti) (VectoBac WG, Valent BioSciences, Libertyville, Illinois, United States) against larvae inside the tyre stacks stored outside and deltamethrin (dilution rate of 2.5 ml/100 ml water) (Aqua K-Othrine, Bayer Environmental Science, Leverkusen, Germany)

against adult mosquitoes. Second, larval control of the surrounding area, a predefined perimeter of 500 m, consists of removing potential larval breeding sites for container-breeding *Aedes* spp., or treating breeding sites with either aqueous spray of Bti (VectoBac WG) or Bti/*Bacillus sphaericus* (Bs) granules (VectoMax FG, Valent BioSciences). Larviciding is being repeated every 2 to 3 weeks until the end of October, which is when ambient conditions for IMS become generally unfavourable. Currently, there is no information of any resistance of this species to the above mentioned biocides.

Discussion

In this article, we report to our best knowledge, the first finding of the species *Ae. flavopictus* outside its area of origin, and the possible association of this finding with the import of used tyres. It is unlikely that the specimen found in the Netherlands was imported from other European countries since, to our best knowledge, the species has not been reported in Europe previously. Preliminary results of inquiries at the company confirms that it regularly imports and stores tyres from Japan. This makes it plausible that there was passive transport of *Ae. flavopictus* eggs through the import of used tyres from the area of origin in Japan.

Ae. flavopictus is widely distributed in Japan and South Korea and comprises three subspecies: *Ae. flavopictus* in the Palearctic region of Japan, *Ae. flavopictus downsi* from the Amami and Okinawa islands, and *Ae. flavopictus miyarai* from Ishigaki Island and Iriomote Island of the Ryukyu archipelago [8]. *Ae. flavopictus* has been described as a rare species in these countries [9], confined to forests and its periphery [10] where females of the species are considered severe day biters [8]. However, the results of Chaves analyses [9] suggest that *Ae. flavopictus* might be antagonistically interacting with *Ae. albopictus* in Japan and spreading because of its ability to cope with the variability of changing environments. This analysis suggests the emergence of this species as a new dominant vector species in communities currently associated with known disease vectors such as *Ae. albopictus*. It is important to note that *Ae. flavopictus* has a similar morphology to that of other *Aedes* species [11], including to that of *Ae. albopictus*, and that both species co-occur, occupying similar breeding sites in Japan [9].

In comparison with *Ae. albopictus*, *Ae. flavopictus* is not known to be a vector of infectious diseases in the field. However, under laboratory conditions, the species has been shown to be a competent vector of dengue [12,13]. Besides dengue virus, *Ae. flavopictus* has been found infected in Japan with *Aedes flavivirus* (AEFV), an insect-specific flavivirus that does not seem to infect vertebrates and that replicates exclusively in mosquito derived cell lines [14].

The international transport of used tyres leads to exotic and potentially invasive mosquito species being

introduced to non-endemic areas where they may have the potential to reproduce and establish. The introduction of IMS with potential to transmit vector-borne diseases are of particular concern. While *Ae. flavopictus* is not known to be a vector of infectious diseases in the field, its detection in Europe is worth noting as it raises awareness of the possibility of new species with different vector competence being introduced via known pathways. Its detection also demonstrates the need for proper identification of morphologically similar mosquito species with different vector competence, e.g. *Ae. albopictus*, *Ae. cretinus* and *Ae. flavopictus*, or *Ae. japonicus* and *Ae. koreicus*, and underlines the importance of active mosquito surveillance systems for timely intervention against exotic mosquitoes with potential future public health impact.

Acknowledgements

We thank Mr Jan Bakker for collecting samples. We thank Dr Luis Fernando Chaves for providing information of the species. We thank Mr Anton van der Sommen for his suggestions regarding the manuscript. We thank Mr Ben van Dijk for collecting company information.

Conflict of interest

None declared.

Authors' contributions

AIJ and AS defined the national surveillance strategy at used tyre companies, and the structure and content of the first draft. BvdV provided the molecular analysis information. RvdB, FJ, MD, EM and JV processed the samples in the laboratory. AIJ wrote the first draft of the manuscript and revised it after comments from all co-authors.

References

1. Beeuwkes J, den Hartog W, Dik M, Scholte E. Surveillance and findings of exotic mosquitoes in used tires in The Netherlands: a methodological approach. *Proc Neth Entomol Soc Meet.* 2011;22. Available from: <https://www.nev.nl/pages/publicaties/proceedings/nummers/22/31-38.pdf>
2. Schaffner F, Angel G, Geoffroy B, Hervy J-P, Rhaïem A, Brunhes J. The mosquitoes of Europe: An identification and training programme. [CD-ROM]. Montpellier: IRD Editions & EID Méditerranée; 2001.
3. Becker N, Petric D, Zgomba M, Boase C, Madon M, Dahl C, et al. Mosquitoes and their control. Heidelberg, Dordrecht, London, New York: Kluwer Academic/Plenum Publishers; 2010.
4. van de Vossenberg BTLH, Ibáñez-Justicia A, Metz-Verschure E, van Veen EJ, Bruil-Dieters ML, Scholte EJ. Real-time PCR Tests in Dutch Exotic Mosquito Surveys; Implementation of *Aedes aegypti* and *Aedes albopictus* Identification Tests, and the Development of Tests for the Identification of *Aedes atropalpus* and *Aedes japonicus japonicus* (Diptera: Culicidae). *J Med Entomol.* 2015;52(3):336-50. <https://doi.org/10.1093/jme/tjv020> PMID: 26334807
5. PM 7/129 (1) DNA barcoding as an identification tool for a number of regulated pests. *Bull OEPP.* 2016;46(3):501-37. <https://doi.org/10.1111/epp.12344>
6. Maekawa Y, Ogawa K, Komagata O, Tsuda Y, Sawabe K. DNA barcoding for molecular identification of Japanese mosquitoes. *Jap J Sanit Zool.* 2016;67(3):183-98. <https://doi.org/10.7601/mez.67.183>
7. Sutherst WR, Maywald G, Kriticos D. CLIMEX version 3. User's Guide. Melbourne: Hearne Scientific Software Pty Ltd; 2007. Available from: <https://www.hearne.software/>

- <getattachment/0343c9d5-999f-4880-b9b2-1c3ee9a08fo8/Climex-User-Guide.aspx>
8. Toma T, Miyagi I, Crabtree MB, Miller BR. Investigation of the *aedes* (*Stegomyia*) *flavopictus* complex (Diptera: Culicidae) in Japan by sequence analysis of the internal transcribed spacers of ribosomal DNA. *J Med Entomol.* 2002;39(3):461-8. <https://doi.org/10.1603/0022-2585-39.3.461> PMID: 12061441
 9. Chaves LF. Globally invasive, withdrawing at home: *Aedes albopictus* and *Aedes japonicus* facing the rise of *Aedes flavopictus*. *Int J Biometeorol.* 2016;60(11):1727-38. <https://doi.org/10.1007/s00484-016-1162-7> PMID: 27039106
 10. Sota T, Mogi M. Interspecific variation in desiccation survival time of *Aedes* (*Stegomyia*) mosquito eggs is correlated with habitat and egg size. *Oecologia.* 1992;90(3):353-8. <https://doi.org/10.1007/BF00317691> PMID: 28313521
 11. Tanaka K, Mizusawa K, Saugstad ES. A Revision of the Adult and Larval Mosquitoes of Japan (including the Ryukyu Archipelago and the Ogasawara Islands) and Korea (Diptera: Culicidae). *Contrib Am Entomol Inst.* 1979;16:1-987. Available from: <http://mosquito-taxonomic-inventory.info/sites/mosquito-taxonomic-inventory.info/files/Tanaka%20et%20al%201979.pdf>
 12. Eshita Y. Experimental studies on the transmission of dengue virus by Japanese mosquitoes. *Teikyo Med J.* 1982;5:17-27.
 13. Srisawat R, Phanitchat T, Komalamisra N, Tamori N, Runtuwene L, Noguchi K, et al. Susceptibility of *Aedes flavopictus* miyurai and *Aedes galloisi* mosquito species in Japan to dengue type 2 virus. *Asian Pac J Trop Biomed.* 2016;6(5):446-50. <https://doi.org/10.1016/j.apjtb.2016.03.003>
 14. Hoshino K, Isawa H, Tsuda Y, Sawabe K, Kobayashi M. Isolation and characterization of a new insect flavivirus from *Aedes albopictus* and *Aedes flavopictus* mosquitoes in Japan. *Virology.* 2009;391(1):119-29. <https://doi.org/10.1016/j.virol.2009.06.025> PMID: 19580982

License, supplementary material and copyright

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0) Licence. You may share and adapt the material, but must give appropriate credit to the source, provide a link to the licence and indicate if changes were made.

Any supplementary material referenced in the article can be found in the online version.

This article is copyright of the authors or their affiliated institutions, 2019.